

ARHGAP18 Polyclonal Antibody

Catalog No: YT0317

Reactivity: Human; Mouse

Applications: WB;IHC;IF;ELISA

Target: ARHGAP18

Gene Name: ARHGAP18

Protein Name: Rho GTPase-activating protein 18

Q8N392

Q8K0Q5

Human Gene Id: 93663

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

RHG18. AA range:517-566

Specificity: ARHGAP18 Polyclonal Antibody detects endogenous levels of ARHGAP18

protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution : WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 75kD

1/4

Background: ARHGAP18 belongs to a family of Rho (see MIM 165390) GTPase-activating

proteins that modulate cell signaling (Potkin et al., 2009 [PubMed

19065146]).[supplied by OMIM, Apr 2010],

Function: function:GTPase activator for the Rho-type GTPases by converting them to an

inactive GDP-bound state., sequence caution: Contaminating sequence. Potential poly-A sequence., similarity: Contains 1 Rho-GAP domain., subunit: Interacts with

MPHOSPH6.,

Subcellular Location:

Cytoplasm.

Expression:

Brain, Mast cell, Melanoma, Uterus,

Sort:

2237

No4:

- 1


Host:

Rabbit


Modifications:

Unmodified


Products Images




Western Blot analysis of various cells using ARHGAP18 Polyclonal Antibody




Western Blot analysis of Jurkat cells using ARHGAP18 Polyclonal Antibody



Western blot analysis of RHG18 Antibody. The lane on the right is blocked with the RHG18 peptide.



Immunohistochemistryt analysis of paraffin-embedded human brain, using RHG18 Antibody. The lane on the right is blocked with the RHG18 peptide.



Western blot analysis of the lysates from HepG2 cells using RHG18 antibody.