

CD70 Polyclonal Antibody

Catalog No: YT0774

Reactivity: Human; Mouse

Applications: WB;ELISA

Target: CD70

Fields: >>Cytokine-cytokine receptor interaction

P32970

O55237

Gene Name: CD70

Protein Name: CD70 antigen

Human Gene ld: 970

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

CD70. AA range:68-117

Specificity: CD70 Polyclonal Antibody detects endogenous levels of CD70 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3



Observed Band: 30kD

Cell Pathway: Cytokine-cytokine receptor interaction;

Background: The protein encoded by this gene is a cytokine that belongs to the tumor

necrosis factor (TNF) ligand family. This cytokine is a ligand for

TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B

lymphocytes. It induces proliferation of costimulated T cells, enhances the

generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin sythesis. [provided by RefSeq, Jul 2008],

Function: function:Cytokine that binds to CD27. Plays a role in T-cell activation. Induces

the proliferation of costimulated T-cells and enhances the generation of cytolytic T-

cells., similarity: Belongs to the tumor necrosis factor family., subunit: Homotrimer.,

Subcellular Location:

Membrane; Single-pass type II membrane protein.

Expression : B-cell, Kidney,

Sort: 3650

No1: 14-0701-85

No2: 14-0701-85


No4:

Host: Rabbit


Modifications: Unmodified

Products Images

2/3



Western Blot analysis of various cells using CD70 Polyclonal Antibody diluted at 1:2000



Western blot analysis of the lysates from Jurkat cells using CD70 antibody.