


## Connexin 43 Polyclonal Antibody

<b>Catalog No :</b>	YT1045
<b>Reactivity :</b>	Human;Mouse;Rat;Monkey
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	Connexin 43
<b>Fields :</b>	>>Gap junction;>>Arrhythmogenic right ventricular cardiomyopathy
<b>Gene Name :</b>	GJA1
<b>Protein Name :</b>	Gap junction alpha-1 protein
<b>Human Gene Id :</b>	2697
<b>Human Swiss Prot No :</b>	P17302
<b>Mouse Gene Id :</b>	14609
<b>Mouse Swiss Prot No :</b>	P23242
<b>Rat Gene Id :</b>	24392
<b>Rat Swiss Prot No :</b>	P08050
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human Connexin 43. AA range:332-381
<b>Specificity :</b>	Connexin 43 Polyclonal Antibody detects endogenous levels of Connexin 43 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200


<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	43kD
<b>Cell Pathway :</b>	Gap junction;Arrhythmogenic right ventricular cardiomyopathy (ARVC);
<b>Background :</b>	This gene is a member of the connexin gene family. The encoded protein is a component of gap junctions, which are composed of arrays of intercellular channels that provide a route for the diffusion of low molecular weight materials from cell to cell. The encoded protein is the major protein of gap junctions in the heart that are thought to have a crucial role in the synchronized contraction of the heart and in embryonic development. A related intronless pseudogene has been mapped to chromosome 5. Mutations in this gene have been associated with oculodentodigital dysplasia, autosomal recessive craniometaphyseal dysplasia and heart malformations. [provided by RefSeq, May 2014],
<b>Function :</b>	caution:PubMed:11741837 reported 2 mutations (Phe-11 and Ala-24) linked to non-syndromic autosomal recessive deafness (DFNBG). These mutations have subsequently been shown (PubMed:12457340) to involve the pseudogene of connexin-43 located on chromosome 5.,caution:PubMed:7715640 reported a mutation Pro-364 linked to congenital heart diseases. This was later shown (PubMed:8873667) to be an artifact.,disease:Defects in GJA1 are a cause of hypoplastic left heart syndrome (HLHS) [MIM:241550]. HLHS refers to the abnormal development of the left-sided cardiac structures, resulting in obstruction to blood flow from the left ventricular outflow tract. In addition, the syndrome includes underdevelopment of the left ventricle, aorta, and aortic arch, as well as mitral atresia or stenosis.,disease:Defects in GJA1 are the cause of autosomal dominant oculodentodigital dysplasia (ODDD) [MIM:164200]; al
<b>Subcellular Location :</b>	Cell membrane ; Multi-pass membrane protein . Cell junction, gap junction . Endoplasmic reticulum . Localizes at the intercalated disk (ICD) in cardiomyocytes and the proper localization at ICD is dependent on TMEM65. .
<b>Expression :</b>	Expressed in the heart and fetal cochlea.
<b>Tag :</b>	hot
<b>Sort :</b>	4419
<b>No4 :</b>	1

**Host :** Rabbit**Modifications :** Unmodified

## Products Images



Western blot analysis of lysates from MCF-7/COS7/Mouse Heart, using Connexin 43 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).