


CtBP1 Polyclonal Antibody


Catalog No :	YT1148
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	CtBP1
Fields :	>>Wnt signaling pathway;>>Notch signaling pathway;>>Pathways in cancer;>>Chronic myeloid leukemia
Gene Name :	CTBP1
Protein Name :	C-terminal-binding protein 1
Human Gene Id :	1487
Human Swiss Prot No :	Q13363
Mouse Gene Id :	13016
Mouse Swiss Prot No :	O88712
Rat Gene Id :	29382
Rat Swiss Prot No :	Q9Z2F5
Immunogen :	The antiserum was produced against synthesized peptide derived from human CtBP1. AA range:388-437
Specificity :	CtBP1 Polyclonal Antibody detects endogenous levels of CtBP1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	48kD
Cell Pathway :	WNT;WNT-T CELLNotch;Pathways in cancer;Chronic myeloid leukemia;
Background :	This gene encodes a protein that binds to the C-terminus of adenovirus E1A proteins. This phosphoprotein is a transcriptional repressor and may play a role during cellular proliferation. This protein and the product of a second closely related gene, CTBP2, can dimerize. Both proteins can also interact with a polycomb group protein complex which participates in regulation of gene expression during development. Alternative splicing of transcripts from this gene results in multiple transcript variants. [provided by RefSeq, Jul 2008],
Function :	cofactor:NAD. Required for efficient interaction with E1A. Cofactor binding induces a conformation change.,function:Involved in controlling the equilibrium between tubular and stacked structures in the Golgi complex (By similarity). Co-repressor targeting diverse transcription regulators such as GLIS2. Has dehydrogenase activity.,PTM:ADP-ribosylated; when cells are exposed to brefeldin-A (BFA).,PTM:Sumoylation on Lys-428 is promoted by the E3 SUMO-protein ligase CBX4.,PTM:The level of phosphorylation appears to be regulated during the cell cycle. Phosphorylated upon DNA damage, probably by ATM or ATR. Phosphorylation by HIPK2 on Ser-422 induces proteasomal degradation.,similarity:Belongs to the D-isomer specific 2-hydroxyacid dehydrogenase family.,subunit:Interacts with the C-terminus of adenovirus E1A protein, ELK3 and CTIP via their consensus motif P-X-[DNS]-L-[STVA]. Can form homodime
Subcellular Location :	Cytoplasm . Nucleus .
Expression :	Expressed in germinal center B-cells.
Sort :	4649
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images




Western Blot analysis of Jurkat cells using CtBP1 Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using CtBP1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat cells, using CtBP1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4° over night