


**Cyclin B1 mouse Monoclonal Antibody(1A5)**

|                              |   |
|------------------------------|---|
| <b>Catalog No :</b>          | YM3650  |
| <b>Reactivity :</b>          | Human;Rat;Mouse   |
| <b>Applications :</b>        | IHC;IF  |
| <b>Target :</b>              | Cyclin B1   |
| <b>Fields :</b>              | >>FoxO signaling pathway;>>Cell cycle;>>Oocyte meiosis;>>p53 signaling pathway;>>Cellular senescence;>>Progesterone-mediated oocyte maturation;>>Human immunodeficiency virus 1 infection |
| <b>Gene Name :</b>           | CCNB1   |
| <b>Protein Name :</b>        | CCNB1   |
| <b>Human Gene Id :</b>       | 891   |
| <b>Human Swiss Prot No :</b> | P14635  |
| <b>Mouse Swiss Prot No :</b> | P24860  |
| <b>Rat Swiss Prot No :</b>   | P30277  |
| <b>Immunogen :</b>           | Synthetic Peptide of Cyclin B1 at AA range of 60-140  |
| <b>Specificity :</b>         | Cyclin B1 protein detects endogenous levels of CCNB1  |
| <b>Formulation :</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Source :</b>              | Monoclonal, Mouse   |
| <b>Dilution :</b>            | IHC 1:100-200. IF 1:50-200  |
| <b>Purification :</b>        | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.  |
| <b>Concentration :</b>       | 1 mg/ml   |

---

|                               |  |
|-------------------------------|--|
| <b>Storage Stability :</b>    | <u>-15°C to -25°C/1 year(Do not lower than -25°C)</u>  |
| <b>Observed Band :</b>        | <u>55kD</u>  |
| <b>Cell Pathway :</b>         | <u>Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;Oocyte meiosis;p53;Progesterone-mediated oocyte maturation;</u>   |
| <b>Background :</b>           | <u>The protein encoded by this gene is a regulatory protein involved in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). Two alternative transcripts have been found, a constitutively expressed transcript and a cell cycle-regulated transcript, that is expressed predominantly during G2/M phase. The different transcripts result from the use of alternate transcription initiation sites. [provided by RefSeq, Jul 2008],</u>  |
| <b>Function :</b>             | <u>developmental stage:Accumulates steadily during G2 and is abruptly destroyed at mitosis.,function:Essential for the control of the cell cycle at the G2/M (mitosis) transition.,PTM:Ubiquitinated by the SCF(NIPA) complex during interphase, leading to its destruction. Not ubiquitinated during G2/M phases.,similarity:Belongs to the cyclin family.,similarity:Belongs to the cyclin family. Cyclin AB subfamily.,subunit:Interacts with the CDC2 protein kinase to form a serine/threonine kinase holoenzyme complex also known as maturation promoting factor (MPF). The cyclin subunit imparts substrate specificity to the complex. Binds HEI10. Interacts with catalytically active RALBP1 and CDC2 during mitosis to form an endocytotic complex during interphase.,</u> |
| <b>Subcellular Location :</b> | <u>Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.</u>   |
| <b>Expression :</b>           | <u>Breast adenocarcinoma,Lung,Placenta,</u>  |
| <b>Sort :</b>                 | <u>4708</u>  |
| <b>No4 :</b>                  | <u>1</u>   |
| <b>Host :</b>                 | <u>Mouse</u>   |
| <b>Modifications :</b>        | <u>Unmodified</u>  |

## Products Images



Immunohistochemical analysis of paraffin-embedded Human Tonsil Tissue using Cyclin B1 Mouse mAb diluted at 1:200.