

## **MEK-7 Polyclonal Antibody**

Catalog No: YT2725

**Reactivity:** Human; Mouse; Rat

**Applications:** WB;IHC

Target: MKK7

**Fields:** >>MAPK signaling pathway;>>ErbB signaling pathway;>>Protein processing in

endoplasmic reticulum;>>Osteoclast differentiation;>>Tight junction;>>Toll-like receptor signaling pathway;>>T cell receptor signaling pathway;>>Fc epsilon RI

signaling pathway;>>TNF signaling pathway;>>Neurotrophin signaling

pathway;>>GnRH signaling pathway;>>Relaxin signaling pathway;>>Alcoholic liver disease;>>Alzheimer disease;>>Huntington disease;>>Pathways of neurodegeneration - multiple diseases;>>Salmonella infection;>>Yersinia

infection;>>Hepatitis B;>>Kaposi sarcoma-associated herpesvirus

infection;>>Epstein-Barr virus infection;>>Human immunodeficiency virus 1 infection;>>Chemical carcinogenesis - reactive oxygen species;>>Lipid and

atherosclerosis:>>Fluid shear stress and atherosclerosis

Gene Name: MAP2K7

**Protein Name:** Dual specificity mitogen-activated protein kinase kinase 7

Human Gene Id: 5609

**Human Swiss Prot** 014733

No:

Mouse Gene ld: 26400

**Mouse Swiss Prot** 

No:

....

**Rat Gene Id :** 363855

Rat Swiss Prot No: Q4KSH7

Immunogen: The antiserum was produced against synthesized peptide derived from human

MAP2K7. AA range:241-290

**Q8CE90** 



**Specificity:** MEK-7 Polyclonal Antibody detects endogenous levels of MEK-7 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500-2000;IHC 1:50-300

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 47kD

**Cell Pathway:** 

MAPK ERK Growth; MAPK G Protein; ErbB HER; Toll Like; T Cell Receptor; F

c epsilon RI;Neurotrophin;GnRH;

**Background:** The protein encoded by this gene is a dual specificity protein kinase that belongs

to the MAP kinase kinase family. This kinase specifically activates MAPK8/JNK1 and MAPK9/JNK2, and this kinase itself is phosphorylated and activated by MAP

kinase kinase kinases including MAP3K1/MEKK1,

MAP3K2/MEKK2, MAP3K3/MEKK5, and MAP4K2/GCK. This kinase is involved in the primary transport of the primary transport of

in the signal transduction mediating the cell responses to proinflammatory cytokines, and environmental stresses. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Jul 2014],

**Function:** catalytic activity:ATP + a protein = ADP + a

phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation by specific MAP kinase kinase kinases such as

MAP3K1/MEKK1, MAP3K3/MEKK3, MAP3K11/MLK3 and

MAP3K12/DLK., function: Stress activated, dual specificity kinase that activates

the JUN kinases MAPK8/JNK1, MAPK9/JNK2 and

MAPK10/JNK3.,PTM:Activated by phosphorylation on Ser/Thr.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase

kinase subfamily., similarity: Contains 1 protein kinase domain., tissue

specificity:Ubiquitous; with highest level of expression in skeletal muscle. Isoform

3 is found at low levels in placenta, fetal liver, and skeletal muscle.,

Subcellular Location:

Nucleus. Cytoplasm.

**Expression:** Ubiquitous; with highest level of expression in skeletal muscle. Isoform 3 is found

at low levels in placenta, fetal liver, and skeletal muscle.

Tag: orthogonal


**Sort :** 9573

**No4**:


**Host:** Rabbit

Modifications: Unmodified


## **Products Images**



Western Blot analysis of 293 cells using MEK-7 Polyclonal Antibody



Western blot analysis of lysates from 293 cells treated with insulin 0.01U/ml 15', using MAP2K7 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).