


Mindin Polyclonal Antibody


| | |
|------------------------------|---|
| Catalog No : | YT2763 |
| Reactivity : | Human;Mouse;Rat;Monkey |
| Applications : | WB;IHC |
| Target : | Mindin |
| Gene Name : | SPON2 |
| Protein Name : | Spondin-2 |
| Human Gene Id : | 10417 |
| Human Swiss Prot No : | Q9BUD6 |
| Mouse Gene Id : | 100689 |
| Mouse Swiss Prot No : | Q8BMS2 |
| Rat Gene Id : | 171569 |
| Rat Swiss Prot No : | Q9WV75 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human SPON2. AA range:71-120 |
| Specificity : | Mindin Polyclonal Antibody detects endogenous levels of Mindin protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500-2000;IHC 1:50-300 |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |

| | |
|-------------------------------|--|
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Observed Band : | 37kD |
| Background : | function:Cell adhesion protein that promote adhesion and outgrowth of hippocampal embryonic neurons. Binds directly to bacteria and their components and functions as an opsonin for macrophage phagocytosis of bacteria. Essential in the initiation of the innate immune response and represents a unique pattern-recognition molecule in the ECM for microbial pathogens.,similarity:Contains 1 spondin domain.,similarity:Contains 1 TSP type-1 domain.,tissue specificity:Expressed in normal lung tissues but not in lung carcinoma cell lines., |
| Function : | function:Cell adhesion protein that promote adhesion and outgrowth of hippocampal embryonic neurons. Binds directly to bacteria and their components and functions as an opsonin for macrophage phagocytosis of bacteria. Essential in the initiation of the innate immune response and represents a unique pattern-recognition molecule in the ECM for microbial pathogens.,similarity:Contains 1 spondin domain.,similarity:Contains 1 TSP type-1 domain.,tissue specificity:Expressed in normal lung tissues but not in lung carcinoma cell lines., |
| Subcellular Location : | Secreted, extracellular space, extracellular matrix . |
| Expression : | Expressed in normal lung tissue but not in lung carcinoma cell lines. |
| Sort : | 9645 |
| No4 : | 1 |
| Host : | Rabbit |
| Modifications : | Unmodified |


Products Images



Western Blot analysis of COS-7 cells using Mindin Polyclonal Antibody



Western blot analysis of the lysates from HUVEC cells using SPON2 antibody.



Immunohistochemical analysis of paraffin-embedded human lung cancer. 1, Antibody was diluted at 1:200 (4° overnight). 2, Tris-EDTA, pH 9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45 min).