

Pax-2 Polyclonal Antibody

Catalog No :	YT3600
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	Pax-2
Gene Name :	PAX2
Protein Name :	Paired box protein Pax-2
Human Gene Id :	5076
Human Swiss Prot No :	Q02962
Mouse Gene Id :	18504
Mouse Swiss Prot No :	P32114
Immunogen :	The antiserum was produced against synthesized peptide derived from human Pax-2. AA range:144-193
Specificity :	Pax-2 Polyclonal Antibody detects endogenous levels of Pax-2 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 42,35kD

Background : PAX2 encodes paired box gene 2, one of many human homologues of the Drosophila melanogaster gene *prd*. The central feature of this transcription factor gene family is the conserved DNA-binding paired box domain. PAX2 is believed to be a target of transcriptional suppression by the tumor suppressor gene WT1. Mutations within PAX2 have been shown to result in optic nerve colobomas and renal hypoplasia. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Dec 2014],

Function : developmental stage:Mainly in fetal kidney and juvenile nephrogenic rests.,disease:Defects in PAX2 are the cause of renal-coloboma syndrome (RCS) [MIM:120330]; also known as papillorenal syndrome or optic nerve coloboma with renal disease. RCS is an autosomal dominant disease characterized by the association of renal hypoplasia, vesicoureteral reflux and dysplasia of the retina and optic disk.,disease:Defects in PAX2 may be responsible for isolated renal hypoplasia as observed in oligomeganephronia (OMN). OMN is a rare congenital and usually sporadic anomaly characterized by bilateral renal hypoplasia, with a reduced number of enlarged nephrons and without urinary tract abnormalities.,function:Probable transcription factor that may have a role in kidney cell differentiation. Has a critical role in the development of the urogenital tract, the eyes, and the CNS.,similarity:Contains 1 paire

Subcellular Location : Nucleus.

Expression : Expressed in primitive cells of the kidney, ureter, eye, ear and central nervous system.


Sort : 11648

No4 : 1


Host : Rabbit

Modifications : Unmodified


Products Images



Western Blot analysis of various cells using Pax-2 Polyclonal Antibody



Immunohistochemistry analysis of Pax-2 antibody in paraffin-embedded human testis tissue.



Western blot analysis of lysate from 293 cells, using Pax-2 antibody.