

PTH/PTHrP-R Polyclonal Antibody

Catalog No: YT3898

Reactivity: Human; Mouse; Rat

Applications: WB;IHC;IF;ELISA

Target: PTH/PTHrP-R

Fields: >>Neuroactive ligand-receptor interaction;>>Parathyroid hormone synthesis,

secretion and action;>>Endocrine and other factor-regulated calcium

reabsorption

Q03431

P41593

Gene Name: PTH1R

Protein Name: Parathyroid hormone/parathyroid hormone-related peptide receptor

Human Gene Id: 5745

Human Swiss Prot

No:

Mouse Gene Id: 19228

Mouse Swiss Prot

No:

Rat Gene ld: 56813

Rat Swiss Prot No: P25961

Immunogen: The antiserum was produced against synthesized peptide derived from human

PTHR1. AA range:145-194

Specificity: PTH/PTHrP-R Polyclonal Antibody detects endogenous levels of PTH/PTHrP-R

protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

1/4



Dilution: WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not

yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 52kD

Cell Pathway: Neuroactive ligand-receptor interaction;

Background: The protein encoded by this gene is a member of the G-protein coupled receptor

family 2. This protein is a receptor for parathyroid hormone (PTH) and for parathyroid hormone-like hormone (PTHLH). The activity of this receptor is

mediated by G proteins which activate adenylyl cyclase and also a

phosphatidylinositol-calcium second messenger system. Defects in this receptor are known to be the cause of Jansen's metaphyseal chondrodysplasia (JMC), chondrodysplasia Blomstrand type (BOCD), as well as enchodromatosis. Two transcript variants encoding the same protein have been found for this gene.

[provided by RefSeg, May 2010],

Function: disease:Defects in PTH1R are a cause of primary failure of tooth eruption (PFE)

[MIM:125350]. PFE is a rare condition that has high penetrance and variable expressivity and in which tooth retention occurs without evidence of any obvious mechanical interference. Instead, malfunction of the eruptive mechanism itself appears to cause nonankylosed permanent teeth to fail to erupt, although the eruption pathway has been cleared by bone resorption., disease: Defects in

PTH1R are the cause of chondrodysplasia Blomstrand type (BOCD)

[MIM:215045]. BOCD is a severe skeletal dysplasia., disease:Defects in PTH1R are the cause of Eiken syndrome [MIM:600002]; also called Eiken skeletal dysplasia or bone modeling defect of hands and feet. Eiken syndrome is a rare familial autosomal recessive skeletal dysplasia. It is characterized by multiple

epiphyseal dysplasia, with extremely retarded ossification, pri

Subcellular Location:

Cell membrane ; Multi-pass membrane protein .

Expression: Expressed in most tissues. Most abundant in kidney, bone and liver.

Tag: orthogonal,hot

Sort: 13153


No4: 1




Host: Rabbit

Modifications: Unmodified


Products Images




Western Blot analysis of COLO cells using PTH/PTHrP-R Polyclonal Antibody diluted at 1:1000



Immunofluorescence analysis of MCF7 cells, using PTHR1 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using PTHR1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat cells, using PTHR1 Antibody. The lane on the right is blocked with the synthesized peptide.