

## **IL-28Ra Polyclonal Antibody**

YT5089 Catalog No:

Reactivity: Human; Mouse

**Applications:** WB;ELISA

Target: IL-28Ra

Fields: >>Cytokine-cytokine receptor interaction;>>JAK-STAT signaling pathway

**Gene Name:** IL28RA

**Protein Name:** Interleukin-28 receptor subunit alpha

Q8CGK5

**Human Gene Id:** 163702

**Human Swiss Prot** 

Q8IU57

No:

**Mouse Swiss Prot** 

No:

Immunogen: Synthesized peptide derived from the Internal region of human IL-28Ra.

**Specificity:** IL-28Ra Polyclonal Antibody detects endogenous levels of IL-28Ra protein.

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Formulation:

Source: Polyclonal, Rabbit, IgG

WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications. **Dilution:** 

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Concentration:** 1 mg/ml

-15°C to -25°C/1 year(Do not lower than -25°C) **Storage Stability:** 

**Observed Band:** 57kD

**Cell Pathway :** Cytokine-cytokine receptor interaction; Jak\_STAT;

**Background:** The protein encoded by this gene belongs to the class II cytokine receptor

family. This protein forms a receptor complex with interleukine 10 receptor, beta (IL10RB). The receptor complex has been shown to interact with three closely related cytokines, including interleukin 28A (IL28A), interleukin 28B (IL28B), and interleukin 29 (IL29). The expression of all three cytokines can be induced by viral infection. The cells overexpressing this protein have been found to have

enhanced responses to IL28A and IL29, but decreased response to IL28B. Three alternatively spliced transcript variants encoding distinct isoforms have been

reported. [provided by RefSeq, Jul 2008],

**Function:** function: The IL28RA/IL10RB dimer is a receptor for IL28A, IL28B and IL29. The

ligand/receptor complex seems to signal through the Jak-STAT pathway.,similarity:Belongs to the type II cytokine receptor

family., similarity: Contains 1 fibronectin type-III domain., subunit: Heterodimer with

IL10RB.,tissue specificity:Widely expressed.,

Subcellular Location :

Membrane; Single-pass type I membrane protein.

**Expression :** Widely expressed.

**Sort :** 8489


No4:

Host: Rabbit

Modifications: Unmodified

## **Products Images**

2/3



Western Blot analysis of extracts from K562 cells, using IL-28Ra Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000