

## K-cadherin Polyclonal Antibody

Catalog No: YT5111

**Reactivity:** Human; Mouse; Rat

**Applications:** WB;IHC;IF;ELISA

Target: K-cadherin

Gene Name: CDH6

Protein Name: Cadherin-6

Human Gene Id: 1004

**Human Swiss Prot** 

No:

Mouse Gene ld: 12563

**Mouse Swiss Prot** 

No:

Rat Gene ld: 25409

Rat Swiss Prot No: P55280

Immunogen: Synthesized peptide derived from K-cadherin . at AA range: 180-260

**Specificity:** K-cadherin Polyclonal Antibody detects endogenous levels of K-cadherin

protein.

P55285

P97326

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution :** WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000.. IF 1:50-200

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Observed Band:

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

88kD

**Background:** This gene encodes a member of the cadherin superfamily. Cadherins are

membrane glycoproteins that mediate homophilic cell-cell adhesion and play critical roles in cell differentiation and morphogenesis. The encoded protein is a

type II cadherin and may play a role in kidney development as well as

endometrium and placenta formation. Decreased expression of this gene may be associated with tumor growth and metastasis. [provided by RefSeq, May 2011],

**Function:** function:Cadherins are calcium dependent cell adhesion proteins. They

preferentially interact with themselves in a homophilic manner in connecting cells;

cadherins may thus contribute to the sorting of heterogeneous cell

types., similarity: Contains 5 cadherin domains., tissue specificity: Highly expressed in brain, cerebellum, and kidney. Lung, pancreas, and gastric mucosa show a weak expression. Also expressed in certain liver and kidney carcinomas..

Subcellular Location:

Cell membrane; Single-pass type I membrane protein.

**Expression:** Highly expressed in brain, cerebellum, and kidney. Lung, pancreas, and gastric

mucosa show a weak expression. Also expressed in certain liver and kidney

carcinomas.


**Sort**: 8846

**No4**: 1

**Host:** Rabbit

Modifications: Unmodified

## **Products Images**



Western Blot analysis of BT474, A549, SHSY5Y, 293 cells using K-cadherin Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000