

## **Glut5 Polyclonal Antibody**

Catalog No: YT5388

**Reactivity:** Human; Rat; Mouse;

**Applications:** WB;ELISA

Target: Glut5

**Fields:** >> Carbohydrate digestion and absorption

P22732

Q9WV38

Gene Name: SLC2A5

**Protein Name:** Solute carrier family 2 facilitated glucose transporter member 5

Human Gene ld: 6518

**Human Swiss Prot** 

No:

**Mouse Swiss Prot** 

No:

Immunogen: The antiserum was produced against synthesized peptide derived from the N-

terminal region of human SLC2A5. AA range:31-80

**Specificity:** Glut5 Polyclonal Antibody detects endogenous levels of Glut5 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3

Observed Band: 55kD

**Background :** The protein encoded by this gene is a fructose transporter responsible for

fructose uptake by the small intestine. The encoded protein also is necessary for

the increase in blood pressure due to high dietary fructose consumption.

[provided by RefSeq, Jun 2016],

Function: function: Cytochalasin B-sensitive carrier. Seems to function primarily as a

fructose transporter.,induction:By forskolin (in Caco-2 cells).,mass spectrometry: PubMed:11840567,similarity:Belongs to the major facilitator superfamily. Sugar

transporter (TC 2.A.1.1) family. Glucose transporter subfamily.,tissue

specificity:Expressed in small intestine, and at much lower levels in kidney,

skeletal muscle, and adipose tissue.,

Subcellular Location:

Apical cell membrane ; Multi-pass membrane protein . Cell membrane ; Multi-pass membrane protein . Cell membrane, sarcolemma . Localized on the apical

membrane of jejunum villi, but also on lateral plasma membranes of the villi.

Transport to the cell membrane is dependent on RAB11A. .

**Expression:** Detected in skeletal muscle, and in jejunum brush border membrane and

basolateral membrane (at protein level) (PubMed:7619085). Expressed in small

intestine, and at much lower levels in kidney, skeletal muscle, and adipose

tissue.

**Sort :** 6642


**No4**:

**Host:** Rabbit


Modifications: Unmodified

## **Products Images**

2/3



Western Blot analysis of K562 cells using Glut5 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from K562 cells, using SLC2A5 Antibody.