


MCM6 Polyclonal Antibody


Catalog No :	YT5454
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC
Target :	MCM6
Fields :	>>DNA replication;>>Cell cycle
Gene Name :	MCM6
Protein Name :	DNA replication licensing factor MCM6
Human Gene Id :	4175
Human Swiss Prot No :	Q14566
Mouse Gene Id :	17219
Mouse Swiss Prot No :	P97311
Rat Swiss Prot No :	Q62724
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human MCM6. AA range:331-380
Specificity :	MCM6 Polyclonal Antibody detects endogenous levels of MCM6 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	90kD
Cell Pathway :	DNA replication;Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;
Background :	<p>The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 7 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. The phosphorylation of the complex by CDC2 kinase reduces the helicase activity, suggesting a role in the regulation of DNA replication. Single nucleotide polymorphisms in the intron regions of this gene are associated with differential transcriptional activation of the promoter of the neighboring lactase gene and, thereby, i</p>
Function :	<p>function:May be involved in the control of a single round of DNA replication during S phase. Binds to chromatin during G1 and detach from it during S phase as if it licenses the chromatin to replicate.,polymorphism:Intronic variations in MCM6 upstream from the LCT gene are associated with adult-type hypolactasia [MIM:223100] leading to lactose intolerance, or with lactase persistence. Lactose intolerance is a normal physiological phenomenon caused by developmental down-regulation of lactase activity during childhood or early adulthood. A non-coding variation in MCM6 affects the transcriptional regulation of the LCT gene resulting in down-regulation of lactase activity. However the majority of Northern Europeans and some African populations have the ability to maintain lactase activity and digest lactose throughout life (lactase persistence).,PTM:Phosphorylated upon DNA damage, probably b</p>
Subcellular Location :	Nucleus . Chromosome . Binds to chromatin during G1 and detach from it during S phase. .
Expression :	Cervix,Epithelium,
Sort :	9468
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images



Western Blot analysis of HeLa cells using MCM6 Polyclonal Antibody . Secondary antibody(catalog#:RS0002) was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western blot analysis of lysate from HeLa cells, using MCM6 Antibody.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).