


ApoC-III Polyclonal Antibody

Catalog No :	YT5550
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IHC;IF;ELISA
Target :	ApoC-III
Fields :	>>PPAR signaling pathway;>>Cholesterol metabolism
Gene Name :	APOC3
Protein Name :	Apolipoprotein C-III
Human Gene Id :	345
Human Swiss Prot No :	P02656
Mouse Swiss Prot No :	P33622
Immunogen :	The antiserum was produced against synthesized peptide derived from the C-terminal region of human APOC3. AA range:46-95
Specificity :	ApoC-III Polyclonal Antibody detects endogenous levels of ApoC-III protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:10000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band :	11kD
Cell Pathway :	PPAR;
Background :	<p>Apolipoprotein C-III is a very low density lipoprotein (VLDL) protein. APOC3 inhibits lipoprotein lipase and hepatic lipase; it is thought to delay catabolism of triglyceride-rich particles. The APOA1, APOC3 and APOA4 genes are closely linked in both rat and human genomes. The A-I and A-IV genes are transcribed from the same strand, while the A-1 and C-III genes are convergently transcribed. An increase in apoC-III levels induces the development of hypertriglyceridemia. [provided by RefSeq, Jul 2008],</p>
Function :	<p>disease:Defects in APOC3 may be a cause of hyperalphalipoproteinemia [MIM:143470]. Affected individuals show high levels of alpha-lipoprotein (high density lipoprotein/HDL).,function:Inhibits lipoprotein lipase and hepatic lipase and decreases the uptake of lymph chylomicrons by hepatic cells. This suggests that it delays the catabolism of triglyceride-rich particles.,PTM:O-linked glycan consists of Gal-GalNAc disaccharide, further modified with up to 3 sialic acid residues.,similarity:Belongs to the apolipoprotein C3 family.,tissue specificity:Constitutes 50% of the protein fraction of VLDL and 2% of that of HDL. Synthesized predominantly in liver and to a lesser degree in intestine.,</p>
Subcellular Location :	Secreted .
Expression :	Liver.
Sort :	2155
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western Blot analysis of 293T-UV cells using ApoC-III Polyclonal Antibody. Antibody was diluted at 1:500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000