

CD53 Polyclonal Antibody

Catalog No: YT5591

Reactivity: Human; Mouse; Rat

Applications: WB;ELISA

Target: CD53

Gene Name: CD53

Protein Name: Leukocyte surface antigen CD53

P19397

Q61451

Human Gene Id: 963

Human Swiss Prot

No:

Mouse Gene ld: 12508

Mouse Swiss Prot

No:

Rat Gene ld: 24251

Rat Swiss Prot No: P24485

Immunogen: The antiserum was produced against synthesized peptide derived from the

Internal region of human CD53. AA range:91-140

Specificity: CD53 Polyclonal Antibody detects endogenous levels of CD53 protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 24kD

Background: The protein encoded by this gene is a member of the transmembrane 4

superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins. It contributes to the transduction of CD2-generated signals in T cells and natural killer cells and has been suggested to play a role in growth regulation. Familial deficiency of this gene has been linked to an immunodeficiency associated with recurrent infectious diseases caused by bacteria, fungi and viruses. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],

Function: function: May be involved in growth regulation in hematopoietic

cells., similarity:Belongs to the tetraspanin (TM4SF) family., tissue specificity:B-cells, monocytes, macrophages, neutrophils, single (CD4 or CD8) positive

thymocytes and peripheral T-cells.,

Subcellular Location:

Cell membrane . Cell junction . Membrane; Multi-pass membrane protein. Concentrates in localized microdomains along the plasma membrane at the

contact sites between cells of fused myotubes. .

Expression : B-cells, monocytes, macrophages, neutrophils, single (CD4 or CD8) positive

thymocytes and peripheral T-cells.


Sort: 3607

No4: 1


Host: Rabbit

Modifications: Unmodified

Products Images



Western Blot analysis of KB cells using CD53 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western Blot analysis of KB cells using CD53 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000