

CD314 Polyclonal Antibody

| | |
|------------------------------|--|
| Catalog No : | YT5944 |
| Reactivity : | Human;Rat;Mouse; |
| Applications : | IHC;IF;ELISA |
| Target : | CD314 |
| Fields : | >>Natural killer cell mediated cytotoxicity;>>Malaria |
| Gene Name : | KLRK1 D12S2489E NKG2D |
| Protein Name : | NKG2-D type II integral membrane protein (Killer cell lectin-like receptor subfamily K member 1) (NK cell receptor D) (NKG2-D-activating NK receptor) (CD antigen CD314) |
| Human Gene Id : | 22914 |
| Human Swiss Prot No : | P26718 |
| Mouse Gene Id : | 27007 |
| Mouse Swiss Prot No : | O54709 |
| Immunogen : | Synthetic peptide from human protein at AA range: 167-216 |
| Specificity : | The antibody detects endogenous CD314 |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | IHC 1:50-200, ELISA 1:10000-20000. IF 1:50-200 |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Cell Pathway : Natural killer cell mediated cytotoxicity;

Background : Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of

Function : alternative products:A number of isoforms are produced,function:Receptor for MICA, MICB, ULBP1, ULBP2, ULBP3 (ULBP2>ULBP1>ULBP3) and ULBP4. Plays a role as a receptor for the recognition of MHC class I HLA-E molecules by NK cells and some cytotoxic T-cells. Involved in the immune surveillance exerted by T- and B-lymphocytes.,miscellaneous:Structurally distinct families of ligands for mouse and human NKG2D receptors have been characterized. They might be orthologs.,online information:NKG-2D,similarity:Contains 1 C-type lectin domain.,subunit:Homodimer. Interacts with DAP10. The interaction with DAP10 is required for NKG2D cell surface expression.,tissue specificity:Natural killer cells. Expressed on essentially all CD56+CD3- NK cells from freshly isolated PBMC. Also detected in gamma-delta cells and CD8+ alpha-beta T-cells. Expressed in interferon-producing killer dendritic cells (IKDCs).

Subcellular Location : Cell membrane ; Single-pass type II membrane protein . Colocalized with HCST on the cell surface.

Expression : Expressed in natural killer (NK) cells, CD8(+) alpha-beta and gamma-delta T-cells. Expressed on essentially all CD56+CD3- NK cells from freshly isolated PBMC. Expressed in interferon-producing killer dendritic cells (IKDCs).


Sort : 3538

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200