


TIF-IA Polyclonal Antibody


Catalog No :	YT6132
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC
Target :	TIF-IA
Gene Name :	RRN3 TIFIA
Protein Name :	RNA polymerase I-specific transcription initiation factor RRN3 (Transcription initiation factor IA) (TIF-IA)
Human Gene Id :	54700
Human Swiss Prot No :	Q9NYV6
Mouse Gene Id :	106298
Mouse Swiss Prot No :	B2RS91
Immunogen :	Synthesized peptide derived from human TIF-IA Polyclonal
Specificity :	This antibody detects endogenous levels of TIF-IA.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band :	70kD
Background :	function:Required for efficient transcription initiation by RNA polymerase I.,similarity:Belongs to the RRN3 family.,
Function :	function:Required for efficient transcription initiation by RNA polymerase I.,similarity:Belongs to the RRN3 family.,
Subcellular Location :	Nucleus, nucleolus .
Expression :	Hippocampus,Kidney,Pooled,Spleen,
Sort :	17157
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western blot analysis of 3T3 lysate, antibody was diluted at 1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4 ° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).