

## DUSP4 Polyclonal Antibody

<b>Catalog No :</b>	YT6141
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	MKP-2
<b>Fields :</b>	>>MAPK signaling pathway
<b>Gene Name :</b>	DUSP4 MKP2 VH2
<b>Protein Name :</b>	Dual specificity protein phosphatase 4 (EC 3.1.3.16) (EC 3.1.3.48) (Dual specificity protein phosphatase hVH2) (Mitogen-activated protein kinase phosphatase 2) (MAP kinase phosphatase 2) (MKP-2)
<b>Human Gene Id :</b>	1846
<b>Human Swiss Prot No :</b>	Q13115
<b>Mouse Gene Id :</b>	319520
<b>Mouse Swiss Prot No :</b>	Q8BFV3
<b>Rat Gene Id :</b>	60587
<b>Rat Swiss Prot No :</b>	Q62767
<b>Immunogen :</b>	Synthesized peptide derived from human DUSP4 Polyclonal
<b>Specificity :</b>	This antibody detects endogenous levels of DUSP4.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000, ELISA 1:10000-20000

<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	44kD
<b>Cell Pathway :</b>	MAPK_ERK_Growth;MAPK_G_Protein;
<b>Background :</b>	<p>The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK1, ERK2 and JNK, is expressed in a variety of tissues, and is localized in the nucleus. Two alternatively spliced transcript variants, encoding distinct isoforms, have been obser</p>
<b>Function :</b>	<p>catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,function:Regulates mitogenic signal transduction by dephosphorylating both Thr and Tyr residues on MAP kinases ERK1 and ERK2.,similarity:Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily.,similarity:Contains 1 rhodanese domain.,similarity:Contains 1 tyrosine-protein phosphatase domain.,</p>
<b>Subcellular Location :</b>	Nucleus .
<b>Expression :</b>	Skin,Uterus,
<b>Tag :</b>	hot
<b>Sort :</b>	5290
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Unmodified

## Products Images

170—  
130—  
100—  
70—  
55—  
40—  
35—  
25—



Western blot analysis of A549 lysate, antibody was diluted at 1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000