

**FoxO4 (phospho Thr451) Polyclonal Antibody**


<b>Catalog No :</b>	YP0382
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	FoxO4
<b>Fields :</b>	>>Ras signaling pathway;>>FoxO signaling pathway;>>Shigellosis
<b>Gene Name :</b>	FOXO4
<b>Protein Name :</b>	Forkhead box protein O4
<b>Human Gene Id :</b>	4303
<b>Human Swiss Prot No :</b>	P98177
<b>Mouse Gene Id :</b>	54601
<b>Mouse Swiss Prot No :</b>	Q9WVH3
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human FOXO4 around the phosphorylation site of Thr451. AA range:417-466
<b>Specificity :</b>	Phospho-FoxO4 (T451) Polyclonal Antibody detects endogenous levels of FoxO4 protein only when phosphorylated at T451.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml

---


<b>Storage Stability :</b>	<u>-15°C to -25°C/1 year(Do not lower than -25°C)</u>
<b>Observed Band :</b>	<u>55kD</u>
<b>Cell Pathway :</b>	<u>Insulin Receptor; B Cell Receptor; Protein_Acetylation</u>
<b>Background :</b>	<u>This gene encodes a member of the O class of winged helix/forkhead transcription factor family. Proteins encoded by this class are regulated by factors involved in growth and differentiation indicating they play a role in these processes. A translocation involving this gene on chromosome X and the homolog of the Drosophila trithorax gene, encoding a DNA binding protein, located on chromosome 11 is associated with leukemia. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2010],</u>
<b>Function :</b>	<u>disease:A chromosomal aberration involving FOXO4 is found in acute leukemias. Translocation t(X;11)(q13;q23) with MLL/HRX. The result is a rogue activator protein.,function:Transcription factor involved in the regulation of the insulin signaling pathway. Binds to insulin-response elements (IREs) and can activate transcription of IGF1. Down-regulates expression of HIF1A and suppresses hypoxia-induced transcriptional activation of HIF1A-modulated genes. Also involved in negative regulation of the cell cycle.,pharmaceutical:A constitutively active FOXO4 mutant where phosphorylation sites Thr-32, Ser-187 and Ser-262 have been mutated to alanine may have therapeutic potential in ERBB2/HER2-overexpressing cancers as it inhibits ERBB2-mediated cell survival, transformation and tumorigenicity.,PTM:Acetylation by CBP, which is induced by peroxidase stress, inhibits transcriptional activity. Dea</u>
<b>Subcellular Location :</b>	<u>Cytoplasm. Nucleus. When phosphorylated, translocated from nucleus to cytoplasm. Dephosphorylation triggers nuclear translocation. Monoubiquitination increases nuclear localization. When deubiquitinated, translocated from nucleus to cytoplasm.</u>
<b>Expression :</b>	<u>Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Isoform zeta is most abundant in the liver, kidney, and pancreas.</u>
<b>Tag :</b>	<u>orthogonal</u>
<b>Sort :</b>	<u>6267</u>
<b>No4 :</b>	<u>1</u>
<b>Host :</b>	<u>Rabbit</u>
<b>Modifications :</b>	<u>Phospho</u>

---

## Products Images



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using FOXO4 (Phospho-Thr451) Antibody



Western blot analysis of lysates from HUVEC cells treated with EGF 200ng/ml 5', using FOXO4 (Phospho-Thr451) Antibody. The lane on the right is blocked with the phospho peptide.