

## c-Myb (phospho Ser12) Polyclonal Antibody

Catalog No: YP0661

**Reactivity:** Human; Mouse

**Applications:** WB;IHC;IF;ELISA

Target: c-Myb

**Fields:** >>PI3K-Akt signaling pathway

Gene Name: MYB

**Protein Name:** Transcriptional activator Myb

P10242

P06876

Human Gene Id: 4602

**Human Swiss Prot** 

ilulliali Swiss

No:

Mouse Gene Id: 17863

**Mouse Swiss Prot** 

No:

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

MYB around the phosphorylation site of Ser12. AA range:1-50

**Specificity:** Phospho-c-Myb (S12) Polyclonal Antibody detects endogenous levels of c-Myb

protein only when phosphorylated at S12.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution :** WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

1/3



Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

**Observed Band:** 65kD

**Cell Pathway:** PI3K/Akt; Protein\_Acetylation

**Background:** This gene encodes a protein with three HTH DNA-binding domains that

functions as a transcription regulator. This protein plays an essential role in the regulation of hematopoiesis. This gene may be aberrently expressed or rearranged or undergo translocation in leukemias and lymphomas, and is considered to be an oncogene. Alternative splicing results in multiple transcript

variants. [provided by RefSeq, Jan 2016],

**Function:** domain: Comprised of 3 domains; an N-terminal DNA-binding domain, a centrally

located transcriptional activation domain and a C-terminal domain involved in transcriptional repression.,function:Transcriptional activator; DNA-binding protein that specifically recognize the sequence 5'-YAAC[GT]G-3'. Plays an important role in the control of proliferation and differentiation of hematopoietic progenitor cells.,PTM:Phosphorylated by NLK on multiple sites, which induces proteasomal

degradation.,PTM:Ubiquitinated; mediated by SIAH1 and leading to its

subsequent proteasomal degradation., similarity: Contains 3 HTH myb-type DNA-binding domains., subunit: Binds MYBBP1A. Interacts with HIPK2, MAF and NLK.,

Subcellular Location:

Nucleus.

**Expression:** Liver, Placenta, Testis,

Tag: orthogonal

**Sort :** 4313


**No4**: 1

Host: Rabbit


Modifications: Phospho

## **Products Images**

2/3



Immunohistochemistry analysis of paraffin-embedded human heart, using MYB (Phospho-Ser12) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HeLa cells treated with Hu 2nM 24h, using MYB (Phospho-Ser12) Antibody. The lane on the right is blocked with the phospho peptide.