


APC1 (phospho Ser355) Polyclonal Antibody


| | |
|------------------------------|--|
| Catalog No : | YP1068 |
| Reactivity : | Human;Mouse |
| Applications : | IHC;IF;ELISA |
| Target : | APC1 |
| Fields : | >>Cell cycle;>>Oocyte meiosis;>>Ubiquitin mediated proteolysis;>>Progesterone-mediated oocyte maturation;>>Human T-cell leukemia virus 1 infection |
| Gene Name : | ANAPC1 |
| Protein Name : | Anaphase-promoting complex subunit 1 |
| Human Gene Id : | 64682 |
| Human Swiss Prot No : | Q9H1A4 |
| Mouse Gene Id : | 17222 |
| Mouse Swiss Prot No : | P53995 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human APC1 around the phosphorylation site of Ser355. AA range:321-370 |
| Specificity : | Phospho-APC1 (S355) Polyclonal Antibody detects endogenous levels of APC1 protein only when phosphorylated at S355. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200 |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |

| | |
|-------------------------------|--|
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Molecularweight : | 217kD |
| Cell Pathway : | Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;Oocyte meiosis;Ubiquitin mediated proteolysis;Progesterone-mediated oocyte maturation; |
| Background : | This gene encodes a subunit of the anaphase-promoting complex. This complex is an E3 ubiquitin ligase that regulates progression through the metaphase to anaphase portion of the cell cycle by ubiquitinating proteins which targets them for degradation. [provided by RefSeq, Dec 2011], |
| Function : | function:Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle.,pathway:Protein modification; protein ubiquitination.,PTM:Phosphorylated. Phosphorylation on Ser-355 occurs specifically during mitosis.,similarity:Belongs to the APC1 family.,similarity:Contains 4 PC repeats.,subunit:The APC/C is composed of at least 11 subunits., |
| Subcellular Location : | nucleoplasm,anaphase-promoting complex,cytosol, |
| Expression : | Colon,Epithelium,Ovary,Placenta, |
| Sort : | 2120 |
| No4 : | 1 |
| Host : | Rabbit |
| Modifications : | Phospho |

Products Images



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using APC1 (Phospho-Ser355) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using APC1 (Phospho-Ser355) Antibody. The picture on the right is blocked with the phospho peptide.