


AATM Monoclonal Antibody


Catalog No :	YM0006
Reactivity :	Human;Mouse;Rat;Monkey
Applications :	WB;IF;ELISA
Target :	AATM
Fields :	>>Arginine biosynthesis;>>Alanine, aspartate and glutamate metabolism;>>Cysteine and methionine metabolism;>>Arginine and proline metabolism;>>Tyrosine metabolism;>>Phenylalanine metabolism;>>Phenylalanine, tyrosine and tryptophan biosynthesis;>>Metabolic pathways;>>Carbon metabolism;>>2-Oxocarboxylic acid metabolism;>>Biosynthesis of amino acids;>>Fat digestion and absorption
Gene Name :	GOT2
Protein Name :	Aspartate aminotransferase, mitochondrial
Human Gene Id :	2806
Human Swiss Prot No :	P00505
Mouse Gene Id :	14719
Mouse Swiss Prot No :	P05202
Rat Gene Id :	25721
Rat Swiss Prot No :	P00507
Immunogen :	Purified recombinant fragment of human AATM expressed in E. Coli.
Specificity :	AATM Monoclonal Antibody detects endogenous levels of AATM protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse

Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	48kD
Cell Pathway :	Alanine; aspartate and glutamate metabolism;Cysteine and methionine metabolism;Arginine and proline metabolism;Tyrosine metabolism;Phenylalanine metabolism;Phenylalanine; tyrosine and tryptophan biosy
P References :	<ol style="list-style-type: none">1. Hepatology. 1998 Apr;27(4):1064-74.2. Cell. 2005 Sep 23;122(6):957-68.3. Psychiatr Genet. 2007 Oct;17(5):314.
Background :	Glutamic-oxaloacetic transaminase is a pyridoxal phosphate-dependent enzyme which exists in cytoplasmic and inner-membrane mitochondrial forms, GOT1 and GOT2, respectively. GOT plays a role in amino acid metabolism and the urea and tricarboxylic acid cycles. The two enzymes are homodimeric and show close homology. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2013],
Function :	catalytic activity:L-aspartate + 2-oxoglutarate = oxaloacetate + L-glutamate.,cofactor:Pyridoxal phosphate.,function:Facilitates cellular uptake of long-chain free fatty acids.,miscellaneous:In eukaryotes there are cytoplasmic, mitochondrial and chloroplastic isozymes.,similarity:Belongs to the class-I pyridoxal-phosphate-dependent aminotransferase family.,subunit:Homodimer.,
Subcellular Location :	Mitochondrion matrix . Cell membrane . Exposure to alcohol promotes translocation to the cell membrane. .
Expression :	Epithelium,Gastric mucosa,Muscle,
Sort :	1569
No4 :	1
Host :	Mouse
Modifications :	Unmodified

Products Images



Western Blot analysis using AATM Monoclonal Antibody against HEK293 (1), PC-12 (2), HL-60 (3), BCBL-1 (4), HepG2 (5) and NIH/3T3 (6) cell lysate.



Immunofluorescence analysis of PC-3 (left) and SK-BR-3 (right) cells using AATM Monoclonal Antibody (green). Red: Actin filaments have been labeled with DY-554 phalloidin. Blue: DRAQ5 fluorescent DNA dye.