

**CD166 Monoclonal Antibody**

<b>Catalog No :</b>	YM0105
<b>Reactivity :</b>	Human
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	CD166
<b>Fields :</b>	>>Cell adhesion molecules
<b>Gene Name :</b>	ALCAM
<b>Protein Name :</b>	CD166 antigen
<b>Human Gene Id :</b>	214
<b>Human Swiss Prot No :</b>	Q13740
<b>Mouse Swiss Prot No :</b>	Q61490
<b>Immunogen :</b>	Purified recombinant fragment of CD166 (aa405-524) expressed in E. Coli.
<b>Specificity :</b>	CD166 Monoclonal Antibody detects endogenous levels of CD166 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
<b>Purification :</b>	Affinity purification
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	65kD
<b>Cell Pathway :</b>	Cell adhesion molecules (CAMs);

**P References :**

1. Prostate. 2003 Jan 1;54(1):34-43.
  2. J Clin Endocrinol Metab. 2003 Jul;88(7):3437-43.
  3. J Cell Sci. 2004 Jun 1;117(Pt 13):2841-52.
- 

**Background :**

This gene encodes activated leukocyte cell adhesion molecule (ALCAM), also known as CD166 (cluster of differentiation 166), which is a member of a subfamily of immunoglobulin receptors with five immunoglobulin-like domains (VVC2C2C2) in the extracellular domain. This protein binds to T-cell differentiation antigene CD6, and is implicated in the processes of cell adhesion and migration. Multiple alternatively spliced transcript variants encoding different isoforms have been found. [provided by RefSeq, Aug 2011],

---

**Function :**

domain:The CD6 binding site is located in the N-terminal Ig-like domain.,function:Cell adhesion molecule that binds to CD6. Involved in neurite extension by neurons via heterophilic and homophilic interactions. May play a role in the binding of T- and B-cells to activated leukocytes, as well as in interactions between cells of the nervous system.,similarity:Contains 2 Ig-like V-type (immunoglobulin-like) domains.,similarity:Contains 3 Ig-like C2-type (immunoglobulin-like) domains.,tissue specificity:Spleen, placenta, liver, and weakly in liver. Expressed by activated T-cells, B-cells, monocytes and thymic epithelial cells. Expressed by neurons in the brain. Restricted expression in tumor cell lines. Preferentially expressed in highly metastasizing melanoma cell lines.,

---

**Subcellular Location :**

Cell membrane ; Single-pass type I membrane protein . Cell projection, axon . Cell projection, dendrite . Detected at the immunological synapse, i.e, at the contact zone between antigen-presenting dendritic cells and T-cells (PubMed:15294938, PubMed:16352806). Colocalizes with CD6 and the TCR/CD3 complex at the immunological synapse (PubMed:15294938). .; [Isoform 3]: Secreted .

---

**Expression :**

Detected on hematopoietic stem cells derived from umbilical cord blood (PubMed:24740813). Detected on lymph vessel endothelial cells, skin and tonsil (PubMed:23169771). Detected on peripheral blood monocytes (PubMed:15048703). Detected on monocyte-derived dendritic cells (at protein level) (PubMed:16352806). Detected at low levels in spleen, placenta, liver (PubMed:9502422). Expressed by activated T-cells, B-cells, monocytes and thymic epithelial cells (PubMed:7760007). Isoform 1 and isoform 3 are detected in vein and artery endothelial cells, astrocytes, keratinocytes and artery smooth muscle cells (PubMed:15496415). Expressed by neurons in the brain. Restricted expression in tumor cell lines. Detected in highly metastasizing melanoma cell lines (PubMed:9502422).

---

**Sort :**3420

---

**No4 :**1


---

**Host :**Mouse

---

**Modifications :** Unmodified

## Products Images



Western Blot analysis using CD166 Monoclonal Antibody against truncated Trx-CD166 recombinant protein (1).