

## GATA-4 Monoclonal Antibody

<b>Catalog No :</b>	YM0299
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	GATA-4
<b>Fields :</b>	>>cGMP-PKG signaling pathway;>>Cellular senescence;>>Tight junction;>>Thyroid hormone signaling pathway
<b>Gene Name :</b>	GATA4
<b>Protein Name :</b>	Transcription factor GATA-4
<b>Human Gene Id :</b>	2626
<b>Human Swiss Prot No :</b>	P43694
<b>Mouse Gene Id :</b>	14463
<b>Mouse Swiss Prot No :</b>	Q08369
<b>Immunogen :</b>	Purified recombinant fragment of human GATA-4 expressed in E. Coli.
<b>Specificity :</b>	GATA-4 Monoclonal Antibody detects endogenous levels of GATA-4 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
<b>Purification :</b>	Affinity purification
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Molecularweight :** 45kD

---

**P References :**  
1. J Mol Cell Cardiol. 2007 Dec;43(6):677-85.  
2. Mol Cell Biol. 2000 Oct;20(20):7550-8.  
3. EMBO J. 1997 Sep 15;16(18):5687-96.

---

**Background :** This gene encodes a member of the GATA family of zinc-finger transcription factors. Members of this family recognize the GATA motif which is present in the promoters of many genes. This protein is thought to regulate genes involved in embryogenesis and in myocardial differentiation and function, and is necessary for normal testicular development. Mutations in this gene have been associated with cardiac septal defects. Additionally, alterations in gene expression have been associated with several cancer types. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2015],

---

**Function :** disease:Defects in GATA4 are the cause of atrial septal defect type 2 (ASD2) [MIM:607941]. ASD2 is a congenital heart malformation characterized by incomplete closure of the wall between the atria resulting in blood flow from the left to the right atria. ASD2 patients show other heart abnormalities including ventricular and atrioventricular septal defects, pulmonary valve thickening or insufficiency of the cardiac valves. ASD2 is not associated with defects in the cardiac conduction system or non-cardiac abnormalities.,function:Transcriptional activator. Binds to the consensus sequence 5'-AGATAG-3'. Acts as a transcriptional activator of ANF in cooperation with NKX2-5.,similarity:Contains 2 GATA-type zinc fingers.,subunit:Interacts with ZNF260 (By similarity). Interacts with the homeobox domain of NKX2-5 through its C-terminal zinc finger. Also interacts with JARID2 which represses its a

---

**Subcellular Location :** Nucleus .

---

**Expression :** Heart,Lung,

---

**Sort :** 6486

---

**No4 :** 1

---

**Host :** Mouse

---


**Modifications :** Unmodified

---

---

**Products Images**

---



Western Blot analysis using GATA-4 Monoclonal Antibody against rat fetal heart (1) and adult heart (2) tissues lysate.