

HSP90β Monoclonal Antibody

Catalog No: YM0342

Reactivity: Human; Mouse; Rat; Monkey

Applications: WB;IHC;IF;FCM;ELISA

Target: HSP90B

Fields: >> Protein processing in endoplasmic reticulum; >> PI3K-Akt signaling

pathway;>>Necroptosis;>>Antigen processing and presentation;>>NOD-like

receptor signaling pathway;>>IL-17 signaling pathway;>>Th17 cell

differentiation;>>Progesterone-mediated oocyte maturation;>>Estrogen signaling

pathway;>>Salmonella infection;>>Pathways in cancer;>>Chemical carcinogenesis - receptor activation;>>Prostate cancer;>>Lipid and

atherosclerosis;>>Fluid shear stress and atherosclerosis

Gene Name: HSP90AB1

Protein Name: Heat shock protein HSP 90-beta

P08238

P11499

Human Gene Id: 3326

Human Swiss Prot

No:

Mouse Gene Id: 15516

Mouse Swiss Prot

No:

Rat Gene Id: 301252

Rat Swiss Prot No: P34058

Immunogen : Purified recombinant fragment of human HSP90β expressed in E. Coli.

Specificity: HSP90β Monoclonal Antibody detects endogenous levels of HSP90β protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution: WB 1:500 - 1:2000. IHC 1:200 - 1:1000. IF 1:200 - 1:1000. Flow cytometry:

1:200 - 1:400. ELISA: 1:10000. Not yet tested in other applications.

Purification : Affinity purification

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 83kD

Cell Pathway: PI3K/Akt; Protein_Acetylation

P References : 1. J Biol Chem. 2009 Dec 18;284(51):35381-9.

2. Int J Biol Macromol. 2009 Oct 1;45(3):310-4.

Background: This gene encodes a member of the heat shock protein 90 family; these proteins

are involved in signal transduction, protein folding and degradation and

morphological evolution. This gene encodes the constitutive form of the cytosolic 90 kDa heat-shock protein and is thought to play a role in gastric apoptosis and

inflammation. Alternative splicing results in multiple transcript variants. Pseudogenes have been identified on multiple chromosomes. [provided by

RefSeq, Dec 2012],

Function: function:Molecular chaperone. Has ATPase activity.,PTM:Phosphorylated upon

DNA damage, probably by ATM or ATR., similarity: Belongs to the heat shock protein 90 family., subcellular location: Identified by mass spectrometry in

melanosome fractions from stage I to stage IV., subunit: Homodimer. Interacts with TP53/p53 (By similarity). Interacts with UNC45A. Binding to UNC45A involves 2

UNC45A monomers per HSP90AB1 dimer.,

Subcellular Cytoplasm . Melanosome . Nucleus . Secreted . Cell membrane . Dynein

axonemal particle. Cell surface. Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). Translocates with BIRC2 from the nucleus to the cytoplasm during differentiation (PubMed:18239673).

Secreted when associated with TGFB1 processed form (LAP)

(PubMed:20599762)...

Expression: Amygdala,Brain cortex,Colon,Colon carcinoma,Embryon

Tag: hot

Location:

Sort: 7945


No4: 1




Host: Mouse

Modifications: Unmodified


Products Images




Western Blot analysis using HSP90β Monoclonal Antibody against Jurkat (1), A431 (2), HeLa (3), A549 (4), HEK293 (5), K562 (6), NIH/3T3 (7), PC-12 (8) and Cos7 (9) cell lysate.



Immunohistochemistry analysis of paraffin-embedded kidney cancer tissues (left) and brain tissues (right) with DAB staining using HSP90 β Monoclonal Antibody.



Immunofluorescence analysis of Hela cells using HSP90 β Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Flow cytometric analysis of Hela cells using HSP90β Monoclonal Antibody (green) and negative control (purple).

