


IL-2 Monoclonal Antibody

Catalog No :	YM0370
Reactivity :	Human
Applications :	WB;ELISA
Target :	IL-2
Fields :	>>Cytokine-cytokine receptor interaction;>>Viral protein interaction with cytokine and cytokine receptor;>>PI3K-Akt signaling pathway;>>C-type lectin receptor signaling pathway;>>JAK-STAT signaling pathway;>>Th1 and Th2 cell differentiation;>>Th17 cell differentiation;>>T cell receptor signaling pathway;>>Intestinal immune network for IgA production;>>Type I diabetes mellitus;>>Yersinia infection;>>Chagas disease;>>Measles;>>Human T-cell leukemia virus 1 infection;>>Coronavirus disease - COVID-19;>>Pathways in cancer;>>Autoimmune thyroid disease;>>Inflammatory bowel disease;>>Allograft rejection;>>Graft-versus-host disease
Gene Name :	IL2
Protein Name :	Interleukin-2
Human Gene Id :	3558
Human Swiss Prot No :	P60568
Mouse Swiss Prot No :	P04351
Immunogen :	Purified recombinant fragment of IL-2 expressed in E. Coli.
Specificity :	IL-2 Monoclonal Antibody detects endogenous levels of IL-2 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications. Affinity purification

Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	18kD
Cell Pathway :	Cytokine-cytokine receptor interaction;Jak_STAT;T_Cell_Receptor;Intestinal immune network for IgA production;Type I diabetes mellitus;Autoimmune thyroid disease;Allograft rejection;Graft-versus-host d
P References :	1. Leukemia. 2008 Dec;22(12):2201-7. 2. J Immunol. 2008 Feb 1;180(3):1490-8.
Background :	The protein encoded by this gene is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in the immune response to antigenic stimuli. [provided by RefSeq, Jul 2008],
Function :	disease:A chromosomal aberration involving IL2 is found in a form of T-cell acute lymphoblastic leukemia (T-ALL). Translocation t(4;16)(q26;p13) with involves TNFRSF17.,function:Produced by T-cells in response to antigenic or mitogenic stimulation, this protein is required for T-cell proliferation and other activities crucial to regulation of the immune response. Can stimulate B-cells, monocytes, lymphokine-activated killer cells, natural killer cells, and glioma cells.,online information:Interleukin-2 entry,online information:The Singapore human mutation and polymorphism database,pharmaceutical:Available under the name Proleukin (Chiron). Used in patients with renal cell carcinoma or metastatic melanoma.,similarity:Belongs to the IL-2 family.,
Subcellular Location :	Secreted.
Expression :	Blood,PCR rescued clones,Placenta,Spleen,
Tag :	hot
Sort :	56
No4 :	1
Host :	Mouse
Modifications :	Unmodified

Products Images



Western Blot analysis using IL-2 Monoclonal Antibody against full-length IL-2 recombinant protein with Trx tag (1) and full-length IL-2-hlgGfc transfected HEK293 cell lysate(2).