

Nkx-3.1 Monoclonal Antibody

Catalog No :	YM0477
Reactivity :	Human
Applications :	WB;IHC;IF;FCM;ELISA
Target :	NKX3.1
Fields :	>>Pathways in cancer;>>Prostate cancer
Gene Name :	NKX3-1
Protein Name :	Homeobox protein Nkx-3.1
Human Gene Id :	4824
Human Swiss Prot No :	Q99801
Mouse Swiss Prot No :	P97436
Immunogen :	Purified recombinant fragment of human Nkx-3.1 expressed in E. Coli.
Specificity :	Nkx-3.1 Monoclonal Antibody detects endogenous levels of Nkx-3.1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500 - 1:2000. IHC 1:200 - 1:1000. Flow cytometry: 1:200 - 1:400. ELISA: 1:10000.. IF 1:50-200
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	26kD

Cell Pathway : Pathways in cancer;Prostate cancer;

P References :

1. Exp Mol Med. 2006 Dec 31;38(6):625-33.
2. Exp Biol Med (Maywood). 2008 Mar;233(3):297-309.
- 3.Mol Biol Rep. 2010 Mar;37(3):1505-12.

Background : This gene encodes a homeobox-containing transcription factor. This transcription factor functions as a negative regulator of epithelial cell growth in prostate tissue. Aberrant expression of this gene is associated with prostate tumor progression. Alternate splicing results in multiple transcript variants of this gene. [provided by RefSeq, Jan 2012],

Function : alternative products:Additional isoforms seem to exist,disease:NKX3-1 has been thought to be one of the target gene of the 8p21 loss of heterozygosity, common in prostate cancer, but neither disruption of the coding region of the gene, nor mutations have been found in prostate cancer.,function:Transcription factor, which binds preferentially the consensus sequence 5'-TAAGT[AG]-3' and can behave as a transcriptional repressor. Could play an important role in regulating proliferation of glandular epithelium and in the formation of ducts in prostate.,induction:By androgens and, in the LNCAP cell line, by estrogens. Androgenic control may be lost in prostate cancer cells during tumor progression from an androgen-dependent to an androgen-independent phase.,similarity:Belongs to the NK-3 homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,subunit:Interacts with serum response f

Subcellular Nucleus .

Location :

Expression : Highly expressed in the prostate and, at a lower level, in the testis.


Sort : 10877

No4 : 1


Host : Mouse

Modifications : Unmodified


Products Images



Western Blot analysis using Nkx-3.1 Monoclonal Antibody against LNCaP (1) cell lysate.



Immunohistochemistry analysis of paraffin-embedded human prostate tissues (A, B) with DAB staining using Nkx-3.1 Monoclonal Antibody.



Flow cytometric analysis of PC-3 cells using Nkx-3.1 Monoclonal Antibody (green) and negative control (purple).