

Skp2 p45 Monoclonal Antibody

Catalog No :	YM0579
Reactivity :	Human
Applications :	WB;ELISA
Target :	Skp2
Fields :	>>FoxO signaling pathway;>>Cell cycle;>>Ubiquitin mediated proteolysis;>>mTOR signaling pathway;>>Epstein-Barr virus infection;>>Pathways in cancer;>>Viral carcinogenesis;>>Small cell lung cancer
Gene Name :	SKP2
Protein Name :	S-phase kinase-associated protein 2
Human Gene Id :	6502
Human Swiss Prot No :	Q13309
Mouse Swiss Prot No :	Q9Z0Z3
Immunogen :	Purified recombinant fragment of Skp2 p45 (aa1-130) expressed in E. Coli.
Specificity :	Skp2 p45 Monoclonal Antibody detects endogenous levels of Skp2 p45 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	48kD

Cell Pathway : Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;Ubiquitin mediated proteolysis;Pathways in cancer;Small cell lung cancer;

P References : 1. Acta Biochim Biophys Sin (Shanghai). 2007 Dec;39(12):999-1007.
2. Clin Cancer Res. 2008 Apr 1;14(7):1966-75.

Background : This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbls class; in addition to an F-box, this protein contains 10 tandem leucine-rich repeats. This protein is an essential element of the cyclin A-CDK2 S-phase kinase. It specifically recognizes phosphorylated cyclin-dependent kinase inhibitor 1B (CDKN1B, also referred to as p27 or KIP1) predominantly in S phase and int

Function : function:Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription. Specifically recognizes phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition. Degradation of CDKN1B/p27kip also requires CKS1. Recognizes target proteins ORC1L, CDT1, RBL2, MLL, CDK9, RAG2, FOXO1A, UBP43, and probably MYC, TOB1 and TAL1. Degradation of TAL1 also requires STUB1. Recognizes CDKN1A in association with CCNE1 or CCNE2 and CDK2.,pathway:Protein modification; protein ubiquitination.,similarity:Contains 1 F-box domain.,similarity:Contains 8 LRR (leucine-rich) repeats.,subunit:Part of the SCF(SKP2) complex consisting of CUL1, RBX1, SKP1 and SKP2. Interacts directly with CUL1 and SK

Subcellular Location : Cytoplasm . Nucleus .

Expression : Epithelium,Liver,Placenta,Prostatic carcinoma,


Sort : 16358

No4 : 1

Host : Mouse

Modifications : Unmodified

Products Images



Western Blot analysis using Skp2 p45 Monoclonal Antibody against truncated Trx-Skp2 p45 recombinant protein (1) and GST-Skp2 p45 (aa1-130) recombinant protein (2).