

PDH-E1 α Monoclonal Antibody


Catalog No :	YM1072
Reactivity :	Human;Mouse;Rat;Bovine;Dog;Pig
Applications :	WB;IF
Target :	PDHA1
Fields :	>>Glycolysis / Gluconeogenesis;>>Citrate cycle (TCA cycle);>>Pyruvate metabolism;>>Metabolic pathways;>>Carbon metabolism;>>HIF-1 signaling pathway;>>Glucagon signaling pathway;>>Central carbon metabolism in cancer;>>Diabetic cardiomyopathy
Gene Name :	PDHA1 ODP A
Protein Name :	Pyruvate dehydrogenase E1 component subunit alpha somatic form mitochondrial
Human Gene Id :	5160
Human Swiss Prot No :	P08559
Mouse Gene Id :	18597
Mouse Swiss Prot No :	P35486
Rat Swiss Prot No :	P26284
Immunogen :	Purified recombinant human PDH-E1 α (C-terminus) protein fragments expressed in E.coli.
Specificity :	PDH-E1 α Monoclonal Antibody detects endogenous levels of PDH-E1 α protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:1000 - 1:2000. IF 1:100 - 1:500. Not yet tested in other applications.

Purification :	Affinity purification
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	43kD
Cell Pathway :	Glycolysis / Gluconeogenesis;Citrate cycle (TCA cycle);Valine; leucine and isoleucine biosynthesis;Pyruvate metabolism;Butanoate metabolism;
Background :	<p>The pyruvate dehydrogenase (PDH) complex is a nuclear-encoded mitochondrial multienzyme complex that catalyzes the overall conversion of pyruvate to acetyl-CoA and CO(2), and provides the primary link between glycolysis and the tricarboxylic acid (TCA) cycle. The PDH complex is composed of multiple copies of three enzymatic components: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and lipoamide dehydrogenase (E3). The E1 enzyme is a heterotetramer of two alpha and two beta subunits. This gene encodes the E1 alpha 1 subunit containing the E1 active site, and plays a key role in the function of the PDH complex. Mutations in this gene are associated with pyruvate dehydrogenase E1-alpha deficiency and X-linked Leigh syndrome. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Mar 2010],</p>
Function :	<p>catalytic activity:Pyruvate + [dihydrolipoyllysine-residue acetyltransferase] lipoyllysine = [dihydrolipoyllysine-residue acetyltransferase] S-acetyldihydrolipoyllysine + CO(2).,cofactor:Thiamine pyrophosphate.,disease:Defects in PDHA1 are a cause of pyruvate decarboxylase E1 component deficiency (PDHE1 deficiency) [MIM:312170]. PDHE1 deficiency is the most common enzyme defect in patients with primary lactic acidosis. It is associated with variable clinical phenotypes ranging from neonatal death to prolonged survival complicated by developmental delay, seizures, ataxia, apnea, and in some cases to an X-linked form of Leigh syndrome (LS) (Leigh encephalomyelopathy).,disease:Defects in PDHA1 are the cause of X-linked Leigh syndrome (LS) [MIM:308930]. LS is an early-onset progressive neurodegenerative disorder with a characteristic neuropathology consisting of focal, bilateral lesions in o</p>
Subcellular Location :	Mitochondrion matrix.
Expression :	Ubiquitous.
Sort :	11774
No4 :	1


Host : Mouse

Modifications : Unmodified

Products Images



Western Blot analysis using PDH-E1 α Monoclonal Antibody against A549, HeLa cell lysate.



Immunofluorescence analysis of HeLa cells using PDH-E1 α Monoclonal Antibody.