


## PA28γ Monoclonal Antibody

<b>Catalog No :</b>	YM1070
<b>Reactivity :</b>	Human;Mouse;Dog;Pig
<b>Applications :</b>	WB;IF
<b>Target :</b>	PA28γ
<b>Fields :</b>	>>Proteasome;>>Antigen processing and presentation;>>Hepatitis C
<b>Gene Name :</b>	PSME3
<b>Protein Name :</b>	Proteasome activator complex subunit 3
<b>Human Gene Id :</b>	10197
<b>Human Swiss Prot No :</b>	P61289
<b>Mouse Gene Id :</b>	19192
<b>Mouse Swiss Prot No :</b>	P61290
<b>Immunogen :</b>	Purified recombinant human PA28γ protein fragments expressed in E.coli.
<b>Specificity :</b>	PA28γ Monoclonal Antibody detects endogenous levels of PA28γ protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	WB 1:1000 - 1:2000. IF 1:100 - 1:500. Not yet tested in other applications.
<b>Purification :</b>	Affinity purification
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

<b>Molecularweight :</b>	30kD
<b>Cell Pathway :</b>	Proteasome;Antigen processing and presentation;
<b>Background :</b>	<p>The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. The immunoproteasome contains an alternate regulator, referred to as the 11S regulator or PA28, that replaces the 19S regulator. Three subunits (alpha, beta and gamma) o</p>
<b>Function :</b>	<p>disease:Sera from patients with systemic lupus erythematosus often contain antibodies that react with the Ki antigen.,domain:The C-terminal sequences affect heptamer stability and proteasome affinity.,function:Subunit of the 11S REG-gamma (also called PA28-gamma) proteasome regulator, a donut-shaped homoheptamer which associates with the proteasome. 11S REG-gamma activates the trypsin-like catalytic subunit of the proteasome but inhibits the chymotrypsin-like and postglutamyl-preferring (PGPH) subunits. Facilitates the MDM2-TP53/p53 interaction which promotes ubiquitination- and MDM2-dependent proteasomal degradation of TP53/p53, limiting its accumulation and resulting in inhibited apoptosis after DNA damage. May also be involved in cell cycle regulation.,induction:Up-regulated in thyroid carcinoma cells.,PTM:Phosphorylated by MAP3K3.,similarity:Belongs to the PA28 family.,subcellular lo</p>
<b>Subcellular Location :</b>	Nucleus . Cytoplasm . Localizes to the cytoplasm during mitosis following nuclear envelope breakdown at this distinct stage of the cell cycle which allows its interaction with MAP3K3 kinase. .
<b>Expression :</b>	B-cell,Embryonic kidney,Fetal brain,Human endometrium carcinoma cell line,L
<b>Sort :</b>	11563
<b>No4 :</b>	1

## Products Images



Western Blot analysis using PA28 $\gamma$  Monoclonal Antibody against HepG2, Jurkat, K562 cell lysate.