


**HptII mouse mAb**

<b>Catalog No :</b>	YM1425
<b>Reactivity :</b>	Transgenic
<b>Applications :</b>	WB
<b>Target :</b>	HptII
<b>Gene Name :</b>	hptii
<b>Human Gene Id :</b>	
<b>Human Swiss Prot No :</b>	
<b>Immunogen :</b>	
<b>Specificity :</b>	Transgenic Only.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	wb dilution 1:1000
<b>Purification :</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	37kD
<b>Sort :</b>	7777
<b>No4 :</b>	1

**Host :** Mouse**Modifications :** Unmodified

## Products Images



Western blot detection of HptII in Transgenic rice (leaf) and Non-transgenic rice (leaf) cell lysates using HptII mouse mAb (1:1000 diluted). Predicted band size: 36,38 kDa. Observed band size: 36,38 kDa. Beta-actin mouse mAb (200068-8F10, 1:1000 diluted) was used as a loading control.