

## EIF4EBP3 mouse mAb

Catalog No: YM1455

Reactivity: Transfected

**Applications:** WB

Target: EIF4EBP3

Gene Name: eif4ebp3

**Human Gene Id:** 8637

**Human Swiss Prot** 

No:

**Mouse Swiss Prot** 

No:

**Immunogen:** Recombinant protein of human EIF4EBP3.

O60516

Q80VV3

**Specificity:** Transfected Only.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**Source:** Monoclonal, Mouse

**Dilution:** wb dilution 1:1000

**Purification:** The antibody was affinity-purified from mouse ascites by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

**Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 11kD

**Background:** eukaryotic translation initiation factor 4E binding protein 3(EIF4EBP3) Homo

sapiens This gene encodes a member of the EIF4EBP family, which consists of proteins that bind to eukaryotic translation initiation factor 4E and regulate its



assembly into EIF4F, the multi-subunit translation initiation factor that recognizes the mRNA cap structure. Read-through transcription from the neighboring upstream gene (MASK or ANKHD1) generates a transcript (MASK-BP3) that encodes a protein comprised of the MASK protein sequence for the majority of the protein and a different C-terminus due to an alternate reading frame for the EIF4EBP3 segments. [provided by RefSeq, Oct 2010],

**Function:** 

function:May play a role as a scaffolding protein that may be associated with the abnormal phenotype of leukemia cells. Isoform 2 may possess an antiapoptotic effect and protect cells during normal cell survival through its regulation of caspases.,function:Regulates eIF4E activity by preventing its assembly into the eIF4F complex.,PTM:Phosphorylated.,similarity:Belongs to the eIF4E-binding protein family.,similarity:Belongs to the mask family.,similarity:Contains 1 KH domain.,similarity:Contains 25 ANK repeats.,subunit:EIF4EBP3 interacts with EIF4E.,subunit:Interacts with PTPN11. Isoform 2 interacts with VPR.,tissue specificity:Expression is highest in skeletal muscle, heart, kidney, and pancreas, whereas there is very little expression in brain and thymus.,tissue specificity:Ubiquitous with high expression in cervix, spleen and brain. Expressed in hematopoietic cells with increased expr

Subcellular Location :

cytoplasm, membrane, eukaryotic translation initiation factor 4F complex,

**Expression:** 

Expression is highest in skeletal muscle, heart, kidney, and pancreas, whereas

there is very little expression in brain and thymus.

**Sort**: 5486


No4: 1

Host: Mouse

Modifications: Unmodified

## **Products Images**

2/3



Western blot detection of EIF4EBP3 in CHO-K1 cell lysate(B) and CHO-K1 transfected by EGFP-EIF4EBP3 fragment(A)cell lysate using EIF4EBP3 mouse mAb (1:1000 diluted).