



Negative Control for mouse IgG Primary Antibody

| | |
|-------------------------------|---|
| Catalog No : | VN0005 |
| Reactivity : | Human;Mouse;Rat;Bovine;Pig;Chick; |
| Applications : | IHC;IP;IF;ELISA |
| Human Swiss Prot No : | / |
| Formulation : | PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA |
| Source : | Mouse, Monoclonal/IgG |
| Dilution : | IHC 1:200-1000. IF 1:100-500. ELISA 1:1000-5000 |
| Purification : | Protein G |
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Subcellular Location : | No staining |
| Tag : | ip |
| Sort : | 1 |
| No4 : | 1 |
| Host : | Mouse |
| Modifications : | Unmodified |

Products Images



GAPDH was immunoprecipitated from SW480 whole cell lysate with anti-GAPDH (PTR2136) antibody. Western blot was performed on the immunoprecipitate using anti-GAPDH antibody, and followed by the HRP-conjugated Goat anti-Mouse IgG light chain antibody. Lane 1 (Input): SW480 whole cell lysate Lane 2 (-): Mouse monoclonal IgG (Negative Control) instead of GAPDH antibody in SW480 whole cell lysate. Lane 3 (+): GAPDH antibody IP in SW480 whole cell lysate



Immunohistochemical analysis of paraffin-embedded human brain tissue using anti-Iba1 mouse mAb (YM4765, left) and mouse IgG, kappa monoclonal control (VN005, right).