

COX IV Monoclonal Antibody(6C8), AbFluor 680 Conjugated

| | |
|------------------------------|--|
| Catalog No : | YM2010 |
| Reactivity : | Human;Rat;Mouse |
| Applications : | WB;IHC;IF; |
| Target : | COX IV |
| Fields : | >>Oxidative phosphorylation;>>Metabolic pathways;>>Cardiac muscle contraction;>>Thermogenesis;>>Non-alcoholic fatty liver disease;>>Alzheimer disease;>>Parkinson disease;>>Amyotrophic lateral sclerosis;>>Huntington disease;>>Prion disease;>>Pathways of neurodegeneration - multiple diseases;>>Chemical carcinogenesis - reactive oxygen species;>>Diabetic cardiomyopathy |
| Gene Name : | COX4I1 |
| Protein Name : | Cytochrome c oxidase subunit 4 isoform 1, mitochondrial |
| Human Gene Id : | 1327 |
| Human Swiss Prot No : | P13073 |
| Specificity : | COX IV Monoclonal Antibody(6C8) AbFluor™ 680 Conjugated specially designed for your Immunofluorescence analysis. |
| Formulation : | Liquid in PBS, pH 7.4, containing 0.02% sodium azide as preservative and 50% Glycerol. |
| Source : | Monoclonal, Mouse IgG1 |
| Dilution : | Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows: IHC 1:50-300, IF 1:200 . |
| Purification : | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. |
| Concentration : | 1mg/ml |

| | |
|-------------------------------|---|
| Storage Stability : | Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezi |
| Cell Pathway : | Oxidative phosphorylation;Cardiac muscle contraction;Alzheimer's disease;Parkinson's disease;Huntington's disease; |
| Background : | Cytochrome c oxidase (COX) is the terminal enzyme of the mitochondrial respiratory chain. It is a multi-subunit enzyme complex that couples the transfer of electrons from cytochrome c to molecular oxygen and contributes to a proton electrochemical gradient across the inner mitochondrial membrane. The complex consists of 13 mitochondrial- and nuclear-encoded subunits. The mitochondrially-encoded subunits perform the electron transfer and proton pumping activities. The functions of the nuclear-encoded subunits are unknown but they may play a role in the regulation and assembly of the complex. This gene encodes the nuclear-encoded subunit IV isoform 1 of the human mitochondrial respiratory chain enzyme. It is located at the 3' of the NOC4 (neighbor of COX4) gene in a head-to-head orientation, and shares a promoter with it. Pseudogenes related to this gene are located on chromosomes |
| Function : | function:This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport.,similarity:Belongs to the cytochrome c oxidase IV family.,tissue specificity:Ubiquitous., |
| Subcellular Location : | Mitochondrion inner membrane ; Single-pass membrane protein . |
| Expression : | Ubiquitous. |
| Sort : | 4461 |
| No4 : | 1 |
| Host : | Mouse |
| Modifications : | Unmodified |
| Conjugate : | AbFluor 680 |

Products Images