


Goat Anti Rabbit IgG (H&L)-AbFluor 594

RS3611			
Rabbit			
Elisa;IF;FCM			
Rabbit IgG (H&L)			
1 mg/ml, liquid in 0.01M Phosphate Buffered Saline, pH 7.2, containing 1% BSA, 50% glycerol, 0.02% Sodium Azide			
Goat			
IF (1:200 - 1:1000), FCM (1:100 - 1:1000), Elisa (Use at an assay dependent concentration)			
The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.			
1mg/mL			
Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezi			
Immunoway secondary antibodies are available conjugated to enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources for your each application. Serum adsorbed secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples.			
hot			
1			
1			
Goat			

Conjugate: AbFluor 594

Products Images



Inhaled Lipid Nanoparticles Alleviate Established Pulmonary Fibrosis Small Yang Yang IF Mouse murine lung epithelial-12 (MLE-12) cell

Alexa	Fluor	350	346/442	Blue
Alexa	Fluor	405	401/421	Blue
Alexa	Fluor	488	496/519	Green
Alexa	Fluor	532	532/553	Yellow
Alexa	Fluor	555	555/565	Yellow
Alexa	Fluor	568	578/603	Red/Orange
Alexa	Fluor	594	590/617	Red/Orange
Alexa	Fluor	633	632/647	Red
Alexa	Fluor	647	650/665	Red
Alexa	Fluor	660	663/690	Near IR
Alexa	Fluor	680	679/702	Near IR
Alexa	Fluor	750	749/775	Near IR
Alexa	Fluor	790	784/814	Near IR

To use the Alexa Fluors with fluorescent imagers, use a spectral line of the blue laser diode for Alexa Fluors 405, a cyan (488 nm) laser for Alexa Fluors 488, a yellow (526 nm) laser for Alexa Fluor 550 or 594, and a red (633 nm) laser for Alexa Fluor 649. The Alexa Fluor 680 and 790 fluors are compatible with laser- and filter-based infrared imaging instruments that emit in the 700 nm, and 800 nm