

Recombinant Human IL18 protein ,C- His Tag

YD2207 Catalog No:

Reactivity: Human

>90% as determined by SDS-PAGE **Purity:**

In Vivo: Please contact with the lab for this information.

Sequence: A DNA sequence encoding the human IL18(Tyr37-Asp193) was fused with the

C-terminal His Tag

Q14116

Human Swiss Prot

No:

Formulation: Supplied as solution form in PBS pH 7.5./ Lyophilized from PBS pH 7.5.

Source: E.coli

Dilution: Reconstitute in sterile water for a stock solution

Concentration: >90% as determined by SDS-PAGE

Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 **Storage Stability:**

to 8 °C for one week. Store at -20 to -80 °C for twelve months from the date of

receipt.

Subcellular

Cytoplasm . Secreted . The precursor is cytosolic (PubMed:33883744). In response to inflammasome-activating signals, cleaved and secreted Location:

(PubMed:33883744). Mature form is secreted and released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744). In contrast, the precursor form is not released, due to the presence of an acidic

region that is proteolytically removed by CASP1 during maturation

(PubMed:33883744). The secretion is dependent on protein unfolding and

facilitated by the cargo receptor TMED10 (PubMed:32272059). .

Sort: 14238

Products Images

1/1