

**Recombinant Human EGFR protein ,C- His tag**

<b>Catalog No :</b>	YD2235
<b>Reactivity :</b>	Human
<b>Purity :</b>	>90% as determined by SDS-PAGE
<b>In Vivo :</b>	
<b>Sequence :</b>	A DNA sequence encoding the human EGFR(Met1-Ser645) was fused with the C-terminal His tag
<b>Human Swiss Prot No :</b>	P00533
<b>Formulation :</b>	Supplied as solution form in PBS pH 7.5./ Lyophilized from PBS pH 7.5.
<b>Source :</b>	Mammalian cells
<b>Dilution :</b>	Reconstitute in sterile water for a stock solution
<b>Concentration :</b>	>90% as determined by SDS-PAGE
<b>Storage Stability :</b>	Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 to 8 °C for one week . Store at -20 to -80 °C for twelve months from the date of receipt.
<b>Subcellular Location :</b>	Cell membrane ; Single-pass type I membrane protein . Endoplasmic reticulum membrane ; Single-pass type I membrane protein. Golgi apparatus membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Endosome . Endosome membrane. Nucleus . In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER (PubMed:20674546, PubMed:17909029). Endocytosed upon activation by ligand (PubMed:2790960, PubMed:17182860, PubMed:27153536, PubMed:17909029). Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF) (PubMed:20551055). .; [Isoform 2]: Secreted.
<b>Expression :</b>	Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.
<b>Sort :</b>	923

## Products Images