

CD10-FC recombinant protein

Catalog No :	YD3024
Reactivity :	Human;
Purity :	>90% as determined by SDS-PAGE
Gene Name :	MME
Protein Name :	Neprilysin (EC 3.4.24.11) (Atriopeptidase) (Common acute lymphocytic leukemia antigen) (CALLA) (Enkephalinase) (Neutral endopeptidase 24.11) (NEP) (Neutral endopeptidase) (Skin fibroblast elastase) (S)
Sequence :	Amino acid:52-750,with FC tag.
Human Gene Id :	4311
Human Swiss Prot No :	P08473
Formulation :	Phosphate-buffered solution
Source :	Mammalian cells
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Function :	Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids (PubMed:15283675, PubMed:6208535, PubMed:6349683, PubMed:8168535). Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond (PubMed:17101991, PubMed:6349683). Catalyzes cleavage of bradykinin, substance P and neurotensin peptides (PubMed:6208535). Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9 (PubMed:15283675, PubMed:6349683). Involved in the degradation of atrial natriuretic factor (ANF) and brain natriuretic factor (BNP(1-32)) (PubMed:16254193, PubMed:2531377, PubMed:2972276). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers (PubMed:20876573).
Subcellular Location :	Cell membrane ; Single-pass type II membrane protein .

Products Images