

WT-1-FC recombinant protein

Catalog No: YD3034

Reactivity: Human;

Purity: >90% as determined by SDS-PAGE

Gene Name: WT1

Protein Name: Wilms tumor protein (WT33)

Sequence: Amino acid:392-438, with FC tag.

P19544

Human Gene Id: 7490

Human Swiss Prot

No:

Formulation: Phosphate-buffered solution

Source: Mammalian cells

Storage Stability: -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

Function: Transcription factor that plays an important role in cellular development and cell

survival (PubMed:7862533). Recognizes and binds to the DNA sequence

5'-GCG(T/G)GGGCG-3' (PubMed:17716689, PubMed:25258363,

PubMed:7862533). Regulates the expression of numerous target genes,

including EPO. Plays an essential role for development of the urogenital system. It has a tumor suppressor as well as an oncogenic role in tumor formation. Function may be isoform-specific: isoforms lacking the KTS motif may act as transcription factors (PubMed:15520190). Isoforms containing the KTS motif may bind mRNA and play a role in mRNA metabolism or splicing (PubMed:16934801). Isoform 1

has lower affinity for DNA, and can bind RNA (PubMed:19123921).

Subcellular Location :

Nucleus . Nucleus, nucleolus. Cytoplasm . Note=Isoforms lacking the KTS motif have a diffuse nuclear location (PubMed:15520190). Shuttles between nucleus

and cytoplasm. .; [Isoform 1]: Nucleus speckle .; [Isoform 4]: Nucleus,

nucleoplasm.

Expression: Expressed in the kidney and a subset of hematopoietic cells.



Products Images