

WT1 Polyclonal Antibody

Catalog No :	YN2274
Reactivity :	Human;Rat;Mouse
Applications :	WB;ELISA
Target :	Wilms' Tumor 1
Fields :	>>Transcriptional misregulation in cancer
Gene Name :	WT1
Protein Name :	Wilms tumor protein (WT33)
Human Gene Id :	7490
Human Swiss Prot No :	P19544
Mouse Swiss Prot No :	P22561
Rat Swiss Prot No :	P49952
Immunogen :	Synthesized peptide derived from human protein . at AA range: 310-390
Specificity :	WT1 Polyclonal Antibody detects endogenous levels of protein.
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 49kD

Background : This gene encodes a transcription factor that contains four zinc-finger motifs at the C-terminus and a proline/glutamine-rich DNA-binding domain at the N-terminus. It has an essential role in the normal development of the urogenital system, and it is mutated in a small subset of patients with Wilms tumor. This gene exhibits complex tissue-specific and polymorphic imprinting pattern, with biallelic, and monoallelic expression from the maternal and paternal alleles in different tissues. Multiple transcript variants have been described. In several variants, there is evidence for the use of a non-AUG (CUG) translation initiation codon upstream of, and in-frame with the first AUG. Authors of PMID:7926762 also provide evidence that WT1 mRNA undergoes RNA editing in human and rat, and that this process is tissue-restricted and developmentally regulated. [provided by RefSeq, Mar 2015],

Function : disease:A chromosomal aberration involving WT1 may be a cause of desmoplastic small round cell tumor (DSRCT). Translocation t(11;22)(p13;q12) with EWSR1.,disease:Defects in WT1 are a cause of hypospadias. Hypospadias is a common malformation in which the urethra opens on the ventral side of the penis. It is considered a complex disorder with both genetic and environmental factors involved in the pathogenesis. Hypospadias can occur alone on an apparently multifactorial basis or as part of syndromes.,disease:Defects in WT1 are a cause of Meacham syndrome [MIM:608978]. Meacham syndrome is a rare sporadically occurring multiple malformation syndrome characterized by male pseudohermaphroditism with abnormal internal female genitalia comprising a uterus and double or septate vagina, complex congenital heart defect and diaphragmatic abnormalities.,disease:Defects in WT1 are a cause of Wilms tum

Subcellular Location : Nucleus . Nucleus, nucleolus. Cytoplasm . Isoforms lacking the KTS motif have a diffuse nuclear location (PubMed:15520190). Shuttles between nucleus and cytoplasm. . ; [Isoform 1]: Nucleus speckle . ; [Isoform 4]: Nucleus, nucleoplasm .

Expression : Expressed in the kidney and a subset of hematopoietic cells.

Sort : 24332

No4 : 1

Host : Rabbit

Modifications : Unmodified

Products Images