

CD137L Polyclonal Antibody

Catalog No :	YT0723
Reactivity :	Human;Mouse
Applications :	WB;IF;ELISA
Target :	CD137L
Fields :	>>Cytokine-cytokine receptor interaction
Gene Name :	TNFSF9
Protein Name :	Tumor necrosis factor ligand superfamily member 9
Human Gene Id :	8744
Human Swiss Prot No :	P41273
Mouse Swiss Prot No :	P41274
Immunogen :	The antiserum was produced against synthesized peptide derived from human TNFSF9. AA range:31-80
Specificity :	CD137L Polyclonal Antibody detects endogenous levels of CD137L protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 23kD

Cell Pathway : Cytokine-cytokine receptor interaction;


Background : The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This transmembrane cytokine is a bidirectional signal transducer that acts as a ligand for TNFRSF9/4-1BB, which is a costimulatory receptor molecule in T lymphocytes. This cytokine and its receptor are involved in the antigen presentation process and in the generation of cytotoxic T cells. The receptor TNFRSF9/4-1BB is absent from resting T lymphocytes but rapidly expressed upon antigenic stimulation. The ligand encoded by this gene, TNFSF9/4-1BBL, has been shown to reactivate anergic T lymphocytes in addition to promoting T lymphocyte proliferation. This cytokine has also been shown to be required for the optimal CD8 responses in CD8 T cells. This cytokine is expressed in carcinoma cell lines, and is thought to be involved in T cell-tumor cell interaction.[provided b

Function : function:Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation-induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages.,similarity:Belongs to the tumor necrosis factor family.,subunit:Homotrimer .,tissue specificity:Expressed in brain, placenta, lung, skeletal muscle and kidney.,


Subcellular Location : Membrane; Single-pass type II membrane protein.

Expression : Expressed in brain, placenta, lung, skeletal muscle and kidney.


Products Images




Western Blot analysis of various cells using CD137L Polyclonal Antibody diluted at 1:1000



Western Blot analysis of HuvEc cells using CD137L Polyclonal Antibody diluted at 1:1000



Immunofluorescence analysis of HeLa cells, using TNFSF9 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HUVEC cells, using TNFSF9 Antibody. The lane on the right is blocked with the synthesized peptide.