


DOR-1 Polyclonal Antibody


Catalog No :	YT1403
Reactivity :	Human;Mouse;Rat;Monkey
Applications :	WB;IHC;IF;ELISA
Target :	DOR-1
Fields :	>>cGMP-PKG signaling pathway;>>Sphingolipid signaling pathway;>>Neuroactive ligand-receptor interaction
Gene Name :	OPRD1
Protein Name :	Delta-type opioid receptor
Human Gene Id :	4985
Human Swiss Prot No :	P41143
Mouse Gene Id :	18386
Mouse Swiss Prot No :	P32300
Rat Gene Id :	24613
Rat Swiss Prot No :	P33533
Immunogen :	The antiserum was produced against synthesized peptide derived from human Opioid Receptor-delta. AA range:323-372
Specificity :	DOR-1 Polyclonal Antibody detects endogenous levels of DOR-1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year (Do not lower than -25°C)
Observed Band :	40kD
Cell Pathway :	Neuroactive ligand-receptor interaction;
Background :	function: Inhibits neurotransmitter release by reducing calcium ion currents and increasing potassium ion conductance. Highly stereoselective. receptor for enkephalins., online information: Delta opioid receptor entry, similarity: Belongs to the G-protein coupled receptor 1 family., subunit: Interacts with GPRASP1.,
Function :	function: Inhibits neurotransmitter release by reducing calcium ion currents and increasing potassium ion conductance. Highly stereoselective. receptor for enkephalins., online information: Delta opioid receptor entry, similarity: Belongs to the G-protein coupled receptor 1 family., subunit: Interacts with GPRASP1.,
Subcellular Location :	Cell membrane ; Multi-pass membrane protein .
Expression :	Detected in oocytes (at protein level). Detected in brain cortex, hypothalamus, hippocampus and olfactory bulb. Detected in oocytes.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Opioid Receptor-delta Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HUVEC and COLO205 cells, using Opioid Receptor-delta Antibody. The lane on the right is blocked with the synthesized peptide.