


DNA-PKCS Polyclonal Antibody


Catalog No :	YT1385
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	DNA-PKCS
Fields :	>>Non-homologous end-joining;>>Cell cycle
Gene Name :	PRKDC
Protein Name :	DNA-dependent protein kinase catalytic subunit
Human Gene Id :	5591
Human Swiss Prot No :	P78527
Mouse Gene Id :	19090
Mouse Swiss Prot No :	P97313
Immunogen :	The antiserum was produced against synthesized peptide derived from human DNA-PK. AA range:4061-4110
Specificity :	DNA-PKCS Polyclonal Antibody detects endogenous levels of DNA-PKCS protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	450kD
Cell Pathway :	Non-homologous end-joining;Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;
Background :	This gene encodes the catalytic subunit of the DNA-dependent protein kinase (DNA-PK). It functions with the Ku70/Ku80 heterodimer protein in DNA double strand break repair and recombination. The protein encoded is a member of the PI3/PI4-kinase family.[provided by RefSeq, Jul 2010],
Function :	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Inhibited by wortmannin. Activity of the enzyme seems to be attenuated by autophosphorylation.,function:Serine/threonine-protein kinase that acts as a molecular sensor for DNA damage. Involved in DNA nonhomologous end joining (NHEJ) required for double-strand break (DSB) repair and V(D)J recombination. Must be bound to DNA to express its catalytic properties. Promotes processing of hairpin DNA structures in V(D)J recombination by activation of the hairpin endonuclease artemis (DCLRE1C). The assembly of the DNA-PK complex at DNA ends is also required for the NHEJ ligation step. Required to protect and align broken ends of DNA. May also act as a scaffold protein to aid the localization of DNA repair proteins to the site of damage. Found at the ends of chromosomes, suggesting a further role in the maintenance of
Subcellular Location :	Nucleus . Nucleus, nucleolus .
Expression :	Brain,Cervix carcinoma,Epithelium,Fetal lung,Placen


Products Images




Immunofluorescence analysis of Caco2 cell. 1,primary Antibody was diluted at 1:100(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - AFluor 488 Secondary antibody (catalog No:RS3211) was diluted at 1:500(room temperature, 50min).




Western Blot analysis of various cells using DNA-PKCS Polyclonal Antibody diluted at 1:2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western Blot analysis of HeLa cells using DNA-PKCS Polyclonal Antibody diluted at 1:2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using DNA-PK Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, using DNA-PK Antibody. The lane on the right is blocked with the synthesized peptide.