


## DOCK 180 Polyclonal Antibody

<b>Catalog No :</b>	YT1391
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	DOCK 180
<b>Fields :</b>	>>Focal adhesion;>>Fc gamma R-mediated phagocytosis;>>Regulation of actin cytoskeleton;>>Bacterial invasion of epithelial cells;>>Shigellosis;>>Yersinia infection
<b>Gene Name :</b>	DOCK1
<b>Protein Name :</b>	Dedicator of cytokinesis protein 1
<b>Human Gene Id :</b>	1793
<b>Human Swiss Prot No :</b>	Q14185
<b>Mouse Gene Id :</b>	330662
<b>Mouse Swiss Prot No :</b>	Q8BUR4
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human DOCK1. AA range:1661-1710
<b>Specificity :</b>	DOCK 180 Polyclonal Antibody detects endogenous levels of DOCK 180 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	215kD
<b>Cell Pathway :</b>	Focal adhesion;Regulates Actin and Cytoskeleton;
<b>Background :</b>	<p>This gene encodes a member of the dedicator of cytokinesis protein family. Dedicator of cytokinesis proteins act as guanine nucleotide exchange factors for small Rho family G proteins. The encoded protein regulates the small GTPase Rac, thereby influencing several biological processes, including phagocytosis and cell migration. Overexpression of this gene has also been associated with certain cancers. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2014],</p>
<b>Function :</b>	<p>domain:The DHR-2 domain is necessary and sufficient for the GEF activity.,function:Involved in cytoskeletal rearrangements required for phagocytosis of apoptotic cells and cell motility. Functions as a guanine nucleotide exchange factor (GEF), which activates Rac Rho small GTPases by exchanging bound GDP for free GTP. Its GEF activity may be enhanced by ELMO1.,similarity:Belongs to the DOCK family.,similarity:Contains 1 DHR-1 (CZH-1) domain.,similarity:Contains 1 DHR-2 (CZH-2) domain.,similarity:Contains 1 SH3 domain.,subcellular location:Recruited to membranes via its interaction with phosphatidylinositol 3,4,5-triphosphate.,subunit:Interacts with the SH3 domains of CRK and NCK2 via multiple sites. Interacts with nucleotide-free RAC1 via its DHR-2 domain. Interacts with ELMO1, ELMO2 and probably ELMO3 via its SH3 domain. Interacts with RAC1 and BAI1.,tissue specificity:Highly expressed</p>
<b>Subcellular Location :</b>	Cytoplasm . Membrane . Recruited to membranes via its interaction with phosphatidylinositol 3,4,5-trisphosphate. .
<b>Expression :</b>	Highly expressed in placenta, lung, kidney, pancreas and ovary. Expressed at intermediate level in thymus, testes and colon.

---

## Products Images



Western blot analysis of lysates from HeLa cells, using DOCK1 Antibody. The lane on the right is blocked with the synthesized peptide.