

## **GALR3 Polyclonal Antibody**

Catalog No: YT1846

**Reactivity:** Human; Rat; Mouse;

**Applications:** WB;IF;ELISA

Target: GALR3

**Fields:** >>Neuroactive ligand-receptor interaction

Gene Name: GALR3

**Protein Name:** Galanin receptor type 3

O60755

O88853

Human Gene Id: 8484

**Human Swiss Prot** 

ilulilali Swiss Fio

No:

**Mouse Swiss Prot** 

No:

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

GALR3. AA range:291-340

**Specificity:** GALR3 Polyclonal Antibody detects endogenous levels of GALR3 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:40000. Not yet tested in other

applications.

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

**Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C)

1/2



Observed Band: 40kD

**Cell Pathway:** Neuroactive ligand-receptor interaction;

**Background:** The neuropeptide galanin modulates a variety of physiologic processes including

cognition/memory, sensory/pain processing, hormone secretion, and feeding behavior. The human galanin receptors are G protein-coupled receptors that functionally couple to their intracellular effector through distinct signaling pathways. GALR3 is found in many tissues and may be expressed as 1.4-, 2.4-,

and 5-kb transcripts [provided by RefSeq, Jul 2008],

**Function:** function:Receptor for the hormone galanin.,similarity:Belongs to the G-protein


coupled receptor 1 family.,

Subcellular Location:


Cell membrane; Multi-pass membrane protein.

**Expression:** Uterus,

## **Products Images**



Immunofluorescence analysis of A549 cells, using GALR3 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night