

Krs-1/2 Polyclonal Antibody

Catalog No :	YT2495
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	Krs-1/2
Fields :	>>MAPK signaling pathway;>>Hippo signaling pathway;>>Hippo signaling pathway - multiple species
Gene Name :	STK3/STK4
Protein Name :	Serine/threonine-protein kinase 3/4
Human Gene Id :	6788/6789
Human Swiss Prot No :	Q13188/Q13043
Mouse Gene Id :	56274/58231
Immunogen :	The antiserum was produced against synthesized peptide derived from human Mst1/2. AA range:149-198
Specificity :	Krs-1/2 Polyclonal Antibody detects endogenous levels of Krs-1/2 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 56kD

Cell Pathway : MAPK_ERK_Growth;MAPK_G_Protein;


Background : serine/threonine kinase 3(STK3) Homo sapiens This gene encodes a serine/threonine protein kinase activated by proapoptotic molecules indicating the encoded protein functions as a growth suppressor. Cleavage of the protein product by caspase removes the inhibitory C-terminal portion. The N-terminal portion is transported to the nucleus where it homodimerizes to form the active kinase which promotes the condensation of chromatin during apoptosis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2012],

Function : catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Inhibited by the C-terminal non-catalytic region. Activated by caspase-cleavage. Full activation also requires homodimerization and autophosphorylation of Thr-180, which are inhibited by the proto-oncogene product RAF1.,function:Stress-activated, pro-apoptotic kinase which, following caspase-cleavage, enters the nucleus and induces chromatin condensation followed by internucleosomal DNA fragmentation. Phosphorylates NKX2-1 (By similarity). Phosphorylates and activates LATS1 and LATS2.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SARAH domain.,subcellular location:The caspase-cleaved form cycles between nucleus and cytoplasm.,subunit:Homodimer; mediated via the coil


Subcellular Location : Cytoplasm . Nucleus . The caspase-cleaved form cycles between nucleus and cytoplasm (PubMed:19525978, PubMed:11278283). Phosphorylation at Thr-117 leads to inhibition of nuclear translocation (PubMed:19525978). .

Expression : Expressed at high levels in adult kidney, skeletal and placenta tissues and at very low levels in adult heart, lung and brain tissues.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Mst1/2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, treated with UV 15', using Mst1/2 Antibody. The lane on the right is blocked with the synthesized peptide.