


mGluR-7 Polyclonal Antibody


Catalog No :	YT2747
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	mGluR-7
Fields :	>>Phospholipase D signaling pathway;>>Neuroactive ligand-receptor interaction;>>Glutamatergic synapse
Gene Name :	GRM7
Protein Name :	Metabotropic glutamate receptor 7
Human Gene Id :	2917
Human Swiss Prot No :	Q14831
Mouse Gene Id :	108073
Mouse Swiss Prot No :	Q68ED2
Rat Gene Id :	81672
Rat Swiss Prot No :	P35400
Immunogen :	The antiserum was produced against synthesized peptide derived from human mGluR7. AA range:866-915
Specificity :	mGluR-7 Polyclonal Antibody detects endogenous levels of mGluR-7 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	100kD
Cell Pathway :	Neuroactive ligand-receptor interaction;
Background :	glutamate metabotropic receptor 7(GRM7) Homo sapiens L-glutamate is the major excitatory neurotransmitter in the central nervous system, and it activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors that have been divided into three groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5, and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3, while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Multiple transcript variants encoding different isoforms have been found
Function :	function:Receptor for glutamate. The activity of this receptor is mediated by a G-protein that inhibits adenylate cyclase activity.,similarity:Belongs to the G-protein coupled receptor 3 family.,subunit:Interacts with PICK1.,tissue specificity:Expressed in many areas of the brain, especially in the cerebral cortex, hippocampus, and cerebellum. Expression of GRM7 isoforms in non-neuronal tissues appears to be restricted to isoform 3 and isoform 4.,
Subcellular Location :	Cell membrane ; Multi-pass membrane protein .
Expression :	Expressed in many areas of the brain, especially in the cerebral cortex, hippocampus, and cerebellum. Expression of GRM7 isoforms in non-neuronal tissues appears to be restricted to isoform 3 and isoform 4.


Products Images




Western Blot analysis of various cells using mGluR-7 Polyclonal Antibody



Western Blot analysis of HuvEc cells using mGluR-7 Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using mGluR7 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HUVEC cells, using mGluR7 Antibody. The lane on the right is blocked with the synthesized peptide.