

## KV9.2 Polyclonal Antibody

<b>Catalog No :</b>	YT2516
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC
<b>Target :</b>	KV9.2
<b>Gene Name :</b>	KCNS2
<b>Protein Name :</b>	Potassium voltage-gated channel subfamily S member 2
<b>Human Gene Id :</b>	3788
<b>Human Swiss Prot No :</b>	Q9ULS6
<b>Mouse Gene Id :</b>	16539
<b>Mouse Swiss Prot No :</b>	O35174
<b>Rat Gene Id :</b>	66022
<b>Rat Swiss Prot No :</b>	Q9ER26
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human KCNS2. AA range:197-246
<b>Specificity :</b>	KV9.2 Polyclonal Antibody detects endogenous levels of KV9.2 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000;IHC 1:50-300
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

**Concentration :** 1 mg/ml

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

**Observed Band :** 54kD


**Background :** domain:The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position.,function:Potassium channel subunit. Modulates channel activity and reduces the ion flow.,similarity:Belongs to the potassium channel family. S subfamily.,subcellular location:May not reach the plasma membrane but remain in an intracellular compartment in the absence of KCNB1.,subunit:Heteromultimer with KCNB1 and with KCNB2. Does not form homomultimers. Might also bind to other channel proteins.,

**Function :** domain:The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position.,function:Potassium channel subunit. Modulates channel activity and reduces the ion flow.,similarity:Belongs to the potassium channel family. S subfamily.,subcellular location:May not reach the plasma membrane but remain in an intracellular compartment in the absence of KCNB1.,subunit:Heteromultimer with KCNB1 and with KCNB2. Does not form homomultimers. Might also bind to other channel proteins.,


**Subcellular Location :** Cell membrane ; Multi-pass membrane protein . May not reach the plasma membrane but remain in an intracellular compartment in the absence of KCNB1 or KCNB2. .

**Expression :** Brain,


## Products Images



Western Blot analysis of various cells using KV9.2 Polyclonal Antibody



Western blot analysis of lysates from HepG2 cells, using KCNS2 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).