

CDX2 Monoclonal Antibody(14H6)

Catalog No :	YM3057
Reactivity :	Human;Mouse;Rat
Applications :	WB;IF;IHC
Target :	CDX2
Fields :	>>Gastric cancer
Gene Name :	CDX2
Protein Name :	Homeobox protein CDX-2
Human Gene Id :	1045
Human Swiss Prot No :	Q99626
Mouse Gene Id :	12591
Mouse Swiss Prot No :	P43241
Immunogen :	Synthetic Peptide of CDX2
Specificity :	The antibody detects endogenous CDX2 proteins.
Formulation :	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Source :	Monoclonal, Mouse
Dilution :	WB 1:1000 IHC 1:200 IF 1:200
Purification :	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 42kD


Background : This gene is a member of the caudal-related homeobox transcription factor gene family. The encoded protein is a major regulator of intestine-specific genes involved in cell growth and differentiation. This protein also plays a role in early embryonic development of the intestinal tract. Aberrant expression of this gene is associated with intestinal inflammation and tumorigenesis. [provided by RefSeq, Jan 2012],

Function : function:Involved in the transcriptional regulation of multiple genes expressed in the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine.,PTM:Phosphorylation of Ser-60 mediates the transactivation capacity.,similarity:Belongs to the Caudal homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,


Subcellular Location : Nucleus .

Expression : Detected in small intestine, colon and pancreas.


Products Images




Helicobacter pylori promotes gastric intestinal metaplasia through activation of IRF3-mediated kynurenine pathway. Wanfu Xu IF,IHC Mouse 1:800 gastric mucosa tissue




Liu, Huiming, et al. "The role of CDX2 in renal tubular lesions during diabetic kidney disease." Aging (Albany NY) 13.5 (2021): 6782.




Liu, Huiming, et al. "The role of CDX2 in renal tubular lesions during diabetic kidney disease." *Aging (Albany NY)* 13.5 (2021): 6782.




Immunofluorescence analysis of Hela cell. 1, Amyloid-β Polyclonal Antibody (green) was diluted at 1:200 (4° overnight). (red) was diluted at 1:200 (4° overnight). 2, Goat Anti Rabbit Alexa Fluor 488 Catalog:RS3211 was diluted at 1:1000 (room temperature, 50min). Goat Anti Mouse Alexa Fluor 594 Catalog:RS3608 was diluted at 1:1000 (room temperature, 50min).




Immunohistochemical analysis of paraffin-embedded Human colon tissue. 1, CDX2 Monoclonal Antibody (14H6) was diluted at 1:200 (4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C, 20min). 3, Secondary antibody was diluted at 1:200 (room temperature, 30min). Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Rat-kidney tissue. 1, CDX2 Monoclonal Antibody(14H6) was diluted at 1:200(4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Mouse-testis tissue. 1, CDX2 Monoclonal Antibody(14H6) was diluted at 1:200(4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.




Immunofluorescence analysis of Mouse-kidney tissue. 1, CDX2 Monoclonal Antibody(14H6)(red) was diluted at 1:200(4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of Rat-spleen tissue. 1, CDX2 Monoclonal Antibody (14H6) (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50 min). 3, Picture B: DAPI (blue) 10 min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Western blot analysis of 1) 293T, 2) Mouse Heart tissue, diluted at 1:2000. cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



IHC staining of human rectal cancer tissue, diluted at 1:200.