


mAChR M1 Polyclonal Antibody


Catalog No :	YT2611
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	mAChR M1
Fields :	>>Calcium signaling pathway;>>cAMP signaling pathway;>>Neuroactive ligand-receptor interaction;>>PI3K-Akt signaling pathway;>>Cholinergic synapse;>>Regulation of actin cytoskeleton;>>Alzheimer disease;>>Pathways of neurodegeneration - multiple diseases
Gene Name :	CHRM1
Protein Name :	Muscarinic acetylcholine receptor M1
Human Gene Id :	1128
Human Swiss Prot No :	P11229
Mouse Gene Id :	12669
Mouse Swiss Prot No :	P12657
Rat Gene Id :	25229
Rat Swiss Prot No :	P08482
Immunogen :	The antiserum was produced against synthesized peptide derived from human CHRM1. AA range:295-344
Specificity :	mAChR M1 Polyclonal Antibody detects endogenous levels of mAChR M1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG

Dilution :	WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	51kD
Cell Pathway :	Calcium;Neuroactive ligand-receptor interaction;Regulates Actin and Cytoskeleton;
Background :	The muscarinic cholinergic receptors belong to a larger family of G protein-coupled receptors. The functional diversity of these receptors is defined by the binding of acetylcholine and includes cellular responses such as adenylate cyclase inhibition, phosphoinositide degeneration, and potassium channel mediation. Muscarinic receptors influence many effects of acetylcholine in the central and peripheral nervous system. The muscarinic cholinergic receptor 1 is involved in mediation of vagally-induced bronchoconstriction and in the acid secretion of the gastrointestinal tract. The gene encoding this receptor is localized to 11q13. [provided by RefSeq, Jul 2008],
Function :	function:The muscarinic acetylcholine receptor mediates various cellular responses, including inhibition of adenylate cyclase, breakdown of phosphoinositides and modulation of potassium channels through the action of G proteins. Primary transducing effect is Pi turnover.,similarity:Belongs to the G-protein coupled receptor 1 family.,subunit:Interacts with GPRASP2.,
Subcellular Location :	Cell membrane; Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein.
Expression :	Amygdala,Brain,

Products Images



Western Blot analysis of HepG2 cells using mAChR M1 Polyclonal Antibody



Western blot analysis of lysates from HepG2 cells, using CHR M1 Antibody. The lane on the right is blocked with the synthesized peptide.